

Design and verify: a new scheme for generating cutting-planes

Santanu S. Dey · Sebastian Pokutta

Received: 21 April 2011 / Accepted: 24 January 2013 / Published online: 15 February 2013
© Springer-Verlag Berlin Heidelberg and Mathematical Optimization Society 2013

Abstract A cutting-plane procedure for integer programming (IP) problems usually involves invoking a black-box procedure (such as the Gomory–Chvátal procedure) to *compute* a cutting-plane. In this paper, we describe an alternative paradigm of using the same cutting-plane black-box. This involves two steps. In the first step, we *design* an inequality $cx \leq d$ where c and d are integral, *independent* of the cutting-plane black-box. In the second step, we *verify* that the designed inequality is a valid inequality by verifying that the set $P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\} \cap \mathbb{Z}^n$ is empty using cutting-planes from the black-box. Here P is the feasible region of the linear-programming relaxation of the IP. We refer to the closure of all cutting-planes that can be verified to be valid using a specific cutting-plane black-box as the *verification closure* of the considered cutting-plane black-box. This paper undertakes a systematic study of properties of verification closures of various cutting-plane black-box procedures.

Keywords Verification scheme · Cutting planes · Integer programming

Mathematics Subject Classification 90C10 · 90C57

Santanu S. Dey was supported by NSF Grant CMMI1149400.

S. S. Dey (✉) · S. Pokutta
H. Milton Stewart School of Industrial and Systems Engineering,
Georgia Institute of Technology, Atlanta, GA 30332, USA
e-mail: santanu.dey@isye.gatech.edu

S. Pokutta
e-mail: sebastian.pokutta@isye.gatech.edu

1 Introduction

Cutting-planes are indispensable for solving Integer Programs (IPs). When using generic cutting-planes (like Gomory–Chvátal or split cuts), often the only guiding principal used is that the incumbent fractional point must be separated. In a way, cutting-planes are generated ‘almost blindly’, where we apply some black-box method to constructively compute valid cutting-planes and hope for the right set of cuts to appear that helps in proving optimality or close significant portion of the integrality gap. One possible approach to improve such a scheme would therefore be if we were somehow able to deliberately design strong cutting-planes that were tailor-made, for example, to prove the optimality of known good candidate solutions. This motivates a different paradigm to generate valid cutting-planes for integer programs: First we design cutting-planes which we believe will be useful without considering their validity. Then, once the cutting-planes are designed, we verify that it is valid.

For $n \in \mathbb{N}$, let $[n] = \{1, \dots, n\}$ and for a rational polytope $P \subseteq \mathbb{R}^n$ denote its *integral hull* by $P_I := \text{conv}(P \cap \mathbb{Z}^n)$, where conv represents convex hull. We now precisely describe the verification scheme (abbreviated as: \mathbb{V} -scheme). Let M be an *admissible* cutting-plane procedure (that is, a valid and ‘reasonable’ cutting-plane system—we will formally define these) and let $M(P)$ be the closure with respect to the family of cutting-planes obtained using M . For example, M could represent split cuts and then $M(P)$ represents the split closure of P . Usually using cutting-planes from a cutting-plane procedure M , implies using valid inequalities for $M(P)$ as cutting-planes. In the \mathbb{V} -scheme, we apply the following procedure: We design or guess the inequality $cx \leq d$ where $(c, d) \in \mathbb{Z}^n \times \mathbb{Z}$. To verify that this inequality is valid for P_I , we apply M to $P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}$ and check whether $M(P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}) = \emptyset$. If $M(P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}) = \emptyset$, then $cx \leq d$ is a valid inequality for P_I . This leads us to the following definition.

Definition 1 We say that the inequality $cx \leq d$ is *verifiable* using a cutting plane operator M for a rational polytope $P \subseteq \mathbb{R}^n$ if $c \in \mathbb{Z}^n$, $d \in \mathbb{Z}$ and $M(P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}) = \emptyset$. \square

We might wonder how much we gain from having to only *verify* that a given inequality $cx \leq d$ is valid for P_I , rather than actually *computing* it. In fact at a first glance, it is not even clear that there would be any difference between computing and verifying. The strength of the verification scheme lies in the following inclusion that can be readily verified for admissible cutting-plane procedures:

$$M(P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}) \subseteq M(P) \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}. \quad (1)$$

The interpretation of this inclusion is that an additional inequality $cx \geq d + 1$ appended to the description of P can provide us with crucial extra information when deriving new cutting-planes by using M that is not available when considering P alone. In other words, (1) can potentially be a strict inclusion such that $M(P \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\}) = \emptyset$ while $M(P) \cap \{x \in \mathbb{R}^n \mid cx \geq d + 1\} \neq \emptyset$. This is equivalent to saying that we can *verify* the validity of $cx \leq d$, however we are not able to *compute* $cx \leq d$. To the best of our knowledge, the only paper discussing

a related idea is [4], but theoretical and computational potential of this approach has not been further investigated.

The set obtained by intersecting all cutting-planes verifiable using M will be called the verification closure (abbreviated as: \mathbb{V} -closure) of M and denoted by $\partial M(P)$, that is,

Definition 2 Let M be a cutting plane operator. Then

$$\partial M(P) := \bigcap_{\substack{(c,d) \in \mathbb{Z}^n \times \mathbb{Z} \\ \text{s.t. } M(P \cap \{x \in \mathbb{R}^n \mid cx \geq d+1\}) = \emptyset}} \{x \in \mathbb{R}^n \mid cx \leq d\}. \tag{2}$$

□

Under mild conditions, (1) implies $\partial M(P) \subseteq M(P)$ for all rational polytopes P . (We formally verify this later.) Since there exist inequalities that can be verified but not computed, this inclusion can be proper. We illustrate this in the next example.

Example 1 Given a rational polytope $P \subseteq \mathbb{R}^n$, recall that the split closure of P is defined as,

$$SC(P) = \bigcap_{(\pi, \pi_0) \in \mathbb{Z}^n \times \mathbb{Z}} \text{conv}((P \cap \{x \in \mathbb{R}^n \mid \pi x \leq \pi_0\}) \cup (P \cap \{x \in \mathbb{R}^n \mid \pi x \geq \pi_0 + 1\})).$$

Let $SC^i(P)$ denote the i th split closure P , that is $SC^i(P) = SC(SC^{i-1}(P))$ and $SC^1(P) := SC(P)$.

Consider the following family of polytopes [3] for $n \in \mathbb{N}$:

$$A_n := \left\{ x \in [0, 1]^n \mid \sum_{i \in I} x_i + \sum_{i \notin I} (1 - x_i) \geq \frac{1}{2} \quad \forall I \subseteq [n] \right\}. \tag{3}$$

Note that $(A_n)_I = \emptyset$ and recall that it takes n rounds of split cuts to establish that A_n is infeasible [7]. For simplicity, consider the instance $P := A_3$. Then $SC^2(A_3) \neq \emptyset$ and $SC^3(A_3) = \emptyset$.

We will show that the \mathbb{V} -split closure of A_3 is the empty set, that is, $\partial SC(A_3) = \emptyset$. We first design the inequality $x_1 + x_2 + x_3 \geq 2$. In order to show that the inequality $x_1 + x_2 + x_3 \geq 2$ is verifiable for $\partial SC(A_3)$ we will establish that $SC(Q) = \emptyset$ where $Q := A_3 \cap \{x \in \mathbb{R}^3 \mid x_1 + x_2 + x_3 \leq 1\}$. It is easy to see that $\max\{x_i \mid x \in Q\} < 1$ for $i \in [3]$ and so we obtain that the split cuts $x_i \leq 0$ for $i \in [3]$ are valid for $SC(Q)$. However, $x_1 + x_2 + x_3 \geq \frac{1}{2}$ is in the description of Q . Thus, $SC(Q) = \emptyset$, and so $x_1 + x_2 + x_3 \geq 2$ can be obtained via the \mathbb{V} -split closure, that is, it is valid for $\partial SC(A_3)$. By symmetry, we also obtain that $\partial SC(A_3) \subseteq \{x \in \mathbb{R}^3 \mid x_1 + x_2 + x_3 \leq 1\}$ and so it follows that $\partial SC(A_3) = \emptyset$. □

We note that rank of A_3 with respect to Gomory–Chvátal (GC) cuts [2, 14], Lift-and-project (LP) cuts [1], and Matrix cone cuts (N_0, N, N_+) [16] is also 3 but the \mathbb{V} -rank is 1 for any of these operators.

Outline and contribution This paper undertakes a systematic study of the strengths and weaknesses of the \mathbb{V} -closures. In Sect. 2, we prove basic properties of the \mathbb{V} -closure. In order to present these results, we first describe general classes of reasonable cutting-planes, the so called *admissible cutting-plane procedures*, a machinery developed in [19]. We prove that ∂M is *almost admissible*, that is the \mathbb{V} -schemes satisfy many important properties that all known classes of admissible cutting-plane procedures including GC cuts, lift-and-project cuts, split cuts (SC), and N, N_0, N_+ cuts satisfy.

In Sect. 3, we show first that \mathbb{V} -schemes have natural inherent strength, that is even if M is an arbitrarily weak admissible cutting-plane procedure, ∂M is at least as strong as the GC and the N_0 closures. We then compare the strength of various regular closures (GC cuts, split cuts, and N_0, N, N_+ cuts) with their \mathbb{V} -versions and with each other. For example, we show that $\partial GC(P) \subseteq SC(P)$ and $\partial N_0(P) \subseteq SC(P)$ for every rational polytope P . The complete list of these results is illustrated in Fig. 1.

In Sect. 4, we present upper and lower bounds on the rank of valid inequalities with respect to the \mathbb{V} -closures for a large class of 0/1 problems. These results show that while the \mathbb{V} -closures are strong compared to the regular closures, they are not unrealistically so.

In Sect. 5, we illustrate the strength of the \mathbb{V} -schemes when applied on specific structured problems. We show that facet-defining inequalities of *monotone polytopes* contained in $[0, 1]^n$ have low rank with respect to any ∂M operator. We show that numerous families of inequalities with high GC, N_0 , or N rank [3, 5, 16] (such as clique inequalities) for the *stable set polytope* have a rank of 1 with respect to any ∂M with M being arbitrarily weak and admissible. We will also show that for the subtour elimination relaxation of the *traveling salesman problem* the rank for ∂M with $M \in \{GC, SC, N_0, N, N_+\}$ is in $\Theta(n)$ where n is the number of nodes, that is the rank is $\Theta(\sqrt{\dim(P)})$ with P being the TSP-polytope. It is well-known that for the case of *rational polytopes in \mathbb{R}^2* the GC rank can be arbitrarily large [2]. In contrast, we establish that the rank of rational polytopes in \mathbb{R}^2 with respect to ∂GC is 1.

An extended abstract of the results in this paper is presented in [11].

2 General properties of the \mathbb{V} -closure

Definition 3 [19] Let M be a cutting plane procedure for binary integer linear programs and let $P := \{x \in [0, 1]^n \mid Ax \leq b\}$ be any rational polytope contained in the 0/1 hypercube. The cutting-plane procedure M is called *admissible* if the following holds:

1. VALIDITY: $P_I \subseteq M(P) \subseteq P$.
2. INCLUSION PRESERVATION: If $P \subseteq Q$, then $M(P) \subseteq M(Q)$ for all polytopes $P, Q \subseteq [0, 1]^n$.
3. HOMOGENEITY: $M(F \cap P) = F \cap M(P)$, for all faces F of $[0, 1]^n$.
4. SINGLE COORDINATE ROUNDING: If $x_i \leq \epsilon < 1$ (or $x_i \geq \epsilon > 0$) is valid for P , then $x_i \leq 0$ (or $x_i \geq 1$) is valid for $M(P)$.
5. COMMUTING WITH COORDINATE FLIPS AND DUPLICATIONS: $\tau_i(M(P)) = M(\tau_i(P))$, where τ_i is either one of the following two operations: (i)Coordinate

- flip: $\tau_i : [0, 1]^n \rightarrow [0, 1]^n$ with $(\tau_i(x))_i = (1 - x_i)$ and $(\tau_i(x))_j = x_j$ for $j \in [n] \setminus \{i\}$; (ii) Coordinate Duplication: $\tau_i : [0, 1]^n \rightarrow [0, 1]^{n+1}$ with $(\tau_i(x))_{n+1} = x_i$ and $(\tau_i(x))_j = x_j$ for $j \in [n]$.
6. SUBSTITUTION INDEPENDENCE: Let $\varphi_F : \mathbb{R}^n \rightarrow \mathbb{R}^d$ be the projection onto the d -dimensional face F of $[0, 1]^n$. Then we require $\varphi_F(M(P \cap F)) = M(\varphi_F(P \cap F))$.
 7. SHORT VERIFICATION: There exists a polynomial p such that for any inequality $cx \leq d$ that is valid for $M(P)$ there is a set $I \subseteq [m]$ with $|I| \leq p(n)$ such that $cx \leq d$ is valid for $M(\{x \in [0, 1]^n \mid a_i x \leq b_i, i \in I\})$.

Let M be a cutting plane procedure for general integer linear programs. Such an operator M is defined as being admissible if (A.) M satisfies (1.)–(7.) when restricted to polytopes contained in $[0, 1]^n$ and (B.) M satisfies

- (a) (1.) for all polytopes $P \subseteq \mathbb{R}^n$, that is, if $P := \{x \in \mathbb{R}^n \mid Ax \leq b\}$ then $P_I \subseteq M(P) \subseteq P$;
- (b) (2.) for any two polytopes P and Q in \mathbb{R}^n , that is if P and Q are rational polytopes satisfying $P \subseteq Q$ then $M(P) \subseteq M(Q)$;
- (c) (7.) for all polytopes $P := \{x \in \mathbb{R}^n \mid a_i x \leq b_i, i \in [m]\}$, that is there exists a polynomial p such that for any inequality $cx \leq d$ that is valid for $M(P)$ there is a set $I \subseteq [m]$ with $|I| \leq p(n)$ such that $cx \leq d$ is valid for $M(\{x \in \mathbb{R}^n \mid a_i x \leq b_i, i \in I\})$.

and satisfies Strong Homogeneity (which replaces Homogeneity)

8. STRONG HOMOGENEITY: If $P \subseteq F^{\leq} := \{x \in \mathbb{R}^n \mid ax \leq b\}$ and $F = \{x \in \mathbb{R}^n \mid ax = b\}$ where $(a, b) \in \mathbb{Z}^n \times \mathbb{Z}$, then $M(F \cap P) = M(P) \cap F$.

In the following, we assume that $M(P)$ is a closed convex set. If M satisfies all required properties for being admissible except (7.), then we say M is *almost admissible*.

We note here that almost all known classes of cutting-plane schemes such as GC cuts, lift-and-project cuts, split cuts, and N, N_0, N_+ are admissible (cf. [19] for more details). Observe that (1) in Sect. 1 follows from inclusion preservation. In the following we will mostly work with admissible cutting-plane procedures, however most results hold more generally (that is, these results hold due to only a subset of the properties of admissible cutting plane operators) and we will indicate in brackets for each result which of the above properties are used. Since we assume to work with valid cutting planes only, Property 1 is added in the bracket for each result.

Also note that whenever M is admissible, then $M(P) \neq P$ whenever $P \subseteq [0, 1]^n$ with $P \neq \emptyset$ and $P_I = \emptyset$ (see [19]). Note that we did not include commutation of M with coordinate permutations, which is merely *renaming of coordinates*, as we assume this to be trivially true.

Remark 1 (Properties of admissible cutting plane for general IP versus binary IP) Requiring strong homogeneity for general IPs leads to a slightly more restricted class than the requirement of homogeneity in the 0/1 case. Our motivation to add this property to the list of properties satisfied by admissible cutting plane operators are: (1) It is an important property that is used for proving various results about the closures of cutting planes (see for example [6,21]) which holds true for all well-known cutting

plane operators for general IPs. (2) Moreover as we show in Theorem 1, this ‘restrictive’ property is inherited by ∂M due to it holding for M . Since in this paper, we study properties of \mathbb{V} -schemes, this is an appropriate property to study.

Note that another typical property one would expect to hold for well-defined cutting-plane procedures for general IPs is invariance under integral translations. While we have not added this to the list of properties of admissible cutting plane operators, it is straightforward to check that this property holds for ∂M , if it holds for M .

Remark 2 (Property 4 and \mathbb{V} -closures) Observe that any \mathbb{V} -scheme ∂M supports property 4 by definition. To see this, let $P \subseteq [0, 1]^n$ be a polytope and let without loss of generality $P \cap \{x \in \mathbb{R}^n \mid x_i = 0\} = \emptyset$. Then in particular $M(P \cap \{x \in \mathbb{R}^n \mid x_i \leq 0\}) = \emptyset$ and thus $\partial M(P) \subseteq \{x \in \mathbb{R}^n \mid x_i = 1\}$.

Remark 3 (Admissible cutting-plane procedures for compact convex sets) The definition of admissible cutting-plane procedures readily generalizes to compact convex sets. Since the set $M(P)$ is convex and compact, this allows us to iterate M , that is given a compact convex set P , the set $M^i(P) := M(M^{i-1}(P))$ is well-defined for $i \in \mathbb{N}$ (where $M^1(P) := M(P)$). Observe that $\partial M(P)$ is a compact convex set by definition and is well-defined for a general compact convex set P if $M(P)$ is defined for compact convex set P . Therefore, the set $(\partial M)^i(P) := \partial M((\partial M)^{i-1}(P))$ is also well-defined for $i \in \mathbb{N}$ (where $(\partial M)^1(P) := \partial M(P)$).

Finally we remark that the assumption of M being applicable for compact convex sets is not necessary for defining iterations of M . Indeed one may alternatively assume that $M(P)$ is a rational polytope. However, this still leaves difficulty with the definition of $(\partial M)^i$. Moreover, this property is not true for well-known cutting plane operators such as the N, N_+ operator. On the other hand, all the well-known operators described above, that is GC, SC, N, N_0, N_+ , are applicable to general compact convex sets. (See [9] for a definition of GC operator for compact convex sets.)

All polytopes are assumed to be rational polytopes in this paper if not stated otherwise. In this case we can confine ourselves to valid inequalities with integral coefficients. We will use e^n to represent the vector of all ones in \mathbb{R}^n . If the dimension of the vector is obvious from context, then we will use e instead of e^n . Recall that $A_n := \{x \in [0, 1]^n \mid \sum_{i \in I} x_i + \sum_{i \notin I} (1 - x_i) \geq \frac{1}{2} \forall I \subseteq [n]\}$; this set is referred often in the rest of the paper. We will use $\{\alpha x \leq \beta\}$ as a shorthand for $\{x \in \mathbb{R}^n \mid \alpha x \leq \beta\}$ whenever the ambient dimension n is understood from context. Let φ_F be the projection onto the face F of $[0, 1]^n$ and $Q = \varphi_F(P \cap F)$. We simplify the notation $Q = \varphi_F(P \cap F)$ as $Q \cong (P \cap F)$. Moreover, instead of the cumbersome notation $\varphi_F(M(P \cap F)) = M(\varphi_F(P \cap F))$ for substitution independence, we will simply say $M(Q) \cong M(P \cap F)$.

Next we present a technical lemma that we require for the main result of this section.

Lemma 1 *Let $Q \subseteq \mathbb{R}^n$ be a compact set contained in the interior of the set $\{\beta x \leq \zeta\}$ with $(\beta, \zeta) \in \mathbb{Z}^n \times \mathbb{Z}$ and let $(\alpha, \eta) \in \mathbb{Z}^n \times \mathbb{Z}$. Then there exists a positive integer τ such that Q is strictly contained in the set $\{(\alpha + \tau\beta)x \leq \eta + \tau\zeta\}$.*

Proof Since Q is a bounded set, $\alpha x \leq \eta + M$ for all $x \in Q$ for some bounded $M \in \mathbb{R}$. Also since Q is contained in the interior of the set $\{\beta x \leq \zeta\}$, there exists an $\epsilon > 0$

such that $\beta x \leq \zeta - \epsilon$ for all $x \in Q$. Therefore for $\tau \in \mathbb{Z}_+$ satisfying $M/\epsilon < \tau$, we obtain that $(\alpha + \tau\beta)x \leq \eta + M + \tau\zeta - \tau\epsilon < \eta + \tau\zeta$ for all $x \in Q$. \square

We next show that ∂M satisfies almost all properties that we observe in most well-known cutting-plane procedures.

Theorem 1 *Let M be an admissible cutting-plane procedure. Then ∂M is almost admissible. In particular,*

1. For 0/1 polytopes, ∂M satisfies properties (1.) to (6.).
2. If M is defined for general polytopes, then ∂M satisfies property (8.).

Proof It is straightforward to verify (1.), (2.), and (4.)–(6.). The non-trivial part is property (8.) (or (3.) respectively). In fact it follows from the original operator M having this property. We will prove (8.); property (3.) in the case of $P \subseteq [0, 1]^n$ follows *mutatis mutandis*.

First observe that $\partial M(P \cap F) \subseteq \partial M(P)$ and $\partial M(P \cap F) \subseteq F$. Therefore, $\partial M(P \cap F) \subseteq \partial M(P) \cap F$. To verify $\partial M(P \cap F) \supseteq \partial M(P) \cap F$, we show that if $\hat{x} \notin \partial M(P \cap F)$, then $\hat{x} \notin \partial M(P) \cap F$. Observe first that if $\hat{x} \notin P \cap F$, then $\hat{x} \notin \partial M(P) \cap F$. Therefore, we assume that $\hat{x} \in P \cap F$. Hence we need to prove that if $\hat{x} \notin \partial M(P \cap F)$ and $\hat{x} \in P \cap F$, then $\hat{x} \notin \partial M(P)$. Since $\hat{x} \notin \partial M(P \cap F)$, there exists $c \in \mathbb{Z}^n$ and $d \in \mathbb{Z}$ such that $c\hat{x} > d$ and $M(P \cap F \cap \{cx \geq d + 1\}) = \emptyset$. By strong homogeneity of M , we obtain

$$M(P \cap \{cx \geq d + 1\}) \cap F = \emptyset. \tag{4}$$

Let $F^\leq = \{ax \leq b\}$ and $F = \{ax = b\}$ with $P \subseteq F^\leq$. Now observe that (4) is equivalent to saying that $M(P \cap \{cx \geq d + 1\})$ is contained in the interior of the set $\{ax \leq b\}$. Therefore by Lemma 1, there exists a $\tau \in \mathbb{Z}_+$ such that $M(P \cap \{cx \geq d + 1\})$ is contained in the interior of $\{(c + \tau a)x \leq d + 1 + \tau b\}$. Equivalently, $M(P \cap \{cx \geq d + 1\}) \cap \{(c + \tau a)x \geq d + 1 + \tau b\} = \emptyset$ which implies

$$M(P \cap \{cx \geq d + 1\}) \cap (P \cap \{(c + \tau a)x \geq d + 1 + \tau b\}) = \emptyset. \tag{5}$$

Since $P \subseteq F^\leq$, we obtain that

$$P \cap \{(c + \tau a)x \geq d + 1 + \tau b\} \subseteq P \cap \{cx \geq d + 1\}. \tag{6}$$

Now using (5), (6) and the inclusion preservation property of M it follows that $M(P \cap \{(c + \tau a)x \geq d + 1 + \tau b\}) = \emptyset$. Thus $(c + \tau a)x \leq d + \tau b$ is a verifiable inequality for $\partial M(P)$. Moreover note that since $\hat{x} \in P \cap F$, we have that $a\hat{x} = b$. Therefore, $(c + \tau a)\hat{x} = c\hat{x} + \tau b > d + \tau b$, where the last inequality follows from the fact that $c\hat{x} > d$. \square

It can be shown that short verification, that is property (7.) of admissible systems follows whenever $\partial M(P)$ is a rational polyhedron. However, we do not need this property for the results in this paper.

3 Strength and comparisons of \mathbb{V} -closures

In this section, we compare various regular closures and their verification counterparts with each other. We first formally define possible relations between admissible closures and the notation we use.

Definition 4 Let L, M be almost admissible. Then

1. L *refines* M , if for all rational polytopes P we have $L(P) \subseteq M(P)$. We write: $L \subseteq M$. It is indicated by empty arrow heads in Fig. 1.
2. L *strictly refines* M , if L refines M and there exists a rational polytope P such that $L(P) \subsetneq M(P)$. We write: $L \subsetneq M$. It is indicated by a filled arrow heads in Fig. 1.
3. L *is incomparable with* M , if there exist rational polytopes P and Q such that $M(P) \subsetneq L(P)$ and $M(Q) \supsetneq L(Q)$. We write: $L \perp M$. It is indicated with an arrow with circle head and tail in Fig. 1.

In each of the above definitions, if either one of L or M is defined only for polytopes $P \subseteq [0, 1]^n$, then we confine the comparison to this class of polytopes. \square

In Sect. 3.1, we will establish the following result.

Theorem 2 Let M be an admissible cutting plane operator. Then

1. $\partial M \subsetneq M$ (via Properties 1, 2, 4, 6, 7).
2. $\partial M \subseteq GC$ and $\partial M \subseteq N_0$ (via Properties 1, 2, 4).

In Sect. 3.2, we will establish the following result.

Theorem 3 Let L and M be admissible cutting plane operators such that $L \subseteq M$. Then $\partial L \subseteq \partial M$. Moreover,

1. $\partial GC \subsetneq SC$.
2. $\partial N_0 \perp \partial GC$.
3. $\partial N_0 \perp SC$.
4. $\partial N \subsetneq \partial N_0$.

Well-known relations between the operators $\{GC, SC, N_0, N, N_+\}$ and those presented in Theorems 2 and 3 are depicted in Fig. 1.

3.1 Strength of ∂M for arbitrary admissible cutting-plane procedures M

In order to show that ∂M refines M , we require the following technical lemma; see [10] for a similar result. We use the notation $\sigma_P(\cdot)$ to refer to the support function of a set P , that is $\sigma_P(c) = \sup\{cx \mid x \in P\}$.

Lemma 2 Let $P, Q \subseteq \mathbb{R}^n$ be compact convex sets. If $\sigma_P(c) \leq \sigma_Q(c)$ for all $c \in \mathbb{Z}^n$, then $P \subseteq Q$.

Proof For a compact convex set T , we have that $T = \bigcap_{c \in \mathbb{Z}^n} \{x \in \mathbb{R}^n \mid cx \leq \sigma_T(c)\}$. See [10] for a proof. If $\hat{x} \in P$, then $c\hat{x} \leq \sigma_P(c)$ for all $c \in \mathbb{Z}^n$. By assumption $\sigma_P(c) \leq \sigma_Q(c)$, we obtain that $c\hat{x} \leq \sigma_Q(c)$ for all $c \in \mathbb{Z}^n$. However since $Q = \bigcap_{c \in \mathbb{Z}^n} \{x \in \mathbb{R}^n \mid cx \leq \sigma_Q(c)\}$, we obtain that $\hat{x} \in Q$. \square




Fig. 1 Direct and ∇ -closures and their relations. M is an arbitrarily weak admissible system. In order to simplify the figure, we have removed the arcs corresponding to $GC \perp N$, $GC \perp N_+$, $SC \perp N$, $SC \perp N_+$. An arc $A \rightarrow B$ states that A is refined by B —the other edges indicate incompatibility

Proposition 1 (Properties 1, 2, 4, 6, 7) *Let M be admissible. Then $\partial M \subsetneq M$.*

Proof We first verify that $\partial M \subseteq M$. Since $M(P) \subseteq P$ and $\partial M(P) \subseteq P$ (this follows from the definition of ∂M and Lemma 2), both $M(P)$ and $\partial M(P)$ are bounded. Moreover since $M(P)$ is closed by definition, and $\partial M(P)$ is defined as the intersection of halfspaces (thus a closed set), we obtain that $M(P)$ and $\partial M(P)$ are both compact convex sets. Thus, by Lemma 2, it is sufficient to compare the support functions of $M(P)$ and $\partial M(P)$ with respect to integer vectors only. Let $\sigma_{M(P)}(c) = d$ for $c \in \mathbb{Z}^n$. We verify that $\sigma_{\partial M(P)}(c) \leq \lfloor d \rfloor$. Observe that, $M(P \cap \{cx \geq \lfloor d \rfloor + 1\}) \subseteq M(P) \cap \{cx \geq \lfloor d \rfloor + 1\}$, where the inclusion follows from the inclusion preservation property of M . However note that since $cx \leq d$ is a valid inequality for $M(P)$, we obtain that $M(P) \cap \{cx \geq \lfloor d \rfloor + 1\} = \emptyset$. Thus, $M(P \cap \{cx \geq \lfloor d \rfloor + 1\}) = \emptyset$ and so $cx \leq \lfloor d \rfloor$ is a valid inequality for $\partial M(P)$. Equivalently we have $\sigma_{\partial M(P)}(c) \leq \lfloor d \rfloor \leq d = \sigma_{M(P)}(c)$, completing the proof.

Now we verify $\partial M \subsetneq M$. Let $n \in \mathbb{N}$ be such that $M(A_n) \neq \emptyset$ and $M(A_{n-1}) = \emptyset$; such an n exists (due to the coordinate rounding property of M we have that $M(A_1) = \emptyset$ and since M satisfies property 7, there exists $t \in \mathbb{N}$ such that $M(A_t) \neq \emptyset$; (see [19])). We claim that $\partial M(A_n) = \emptyset$ which implies that $\partial M \subsetneq M$ follows.

In order to establish the claim, observe that $M(A_n \cap \{x_n \leq 0\}) \cong M(A_{n-1}) = \emptyset$, where the last equality is due to the choice of n . Therefore $x_n \geq 1$ is valid for $\partial M(A_n)$. Similarly, we can derive the validity of $x_n \leq 0$ for $\partial M(A_n)$. We therefore conclude that $\partial M(A_n) = \emptyset$. □

Remark 4 Recall that ∂M is defined via integral inequalities and $M(P)$ does neither have to be polyhedral nor rational in general. We use compactness and Lemma 2 to confine ourselves to integral normals which allows for the comparison of M and ∂M .

Also observe that for strict inclusion it suffices that there exists a polytope $P \subseteq [0, 1]^n$ with $P_I = \emptyset$ and $M(P) \neq \emptyset$. Then it can be verified that there is a face F of $[0, 1]^n$ (F can be $[0, 1]^n$) such that $M(P \cap F) \neq \emptyset$ but $\partial M(P \cap F) = \emptyset$ via properties 1, 2, and 3 of an admissible operator M . The requirement of P such that $M(P) \neq \emptyset$ and $P_I = \emptyset$, can be considered a weakening of property 7 which is sufficient for the existence of such a polytope.

We next show that even if M is chosen arbitrarily, ∂M is at least as strong as the GC closure and the N_0 closure. Let $M \circ L$ denote the composition of two operators, that is $(M \circ L)(P) = M(L(P))$. Note that here we assume that M is an admissible cutting plane operator applicable to general compact convex sets if $L(P)$ is not polyhedral. Also note that $M \circ L$ is an admissible operator. See [19] for a proof of this for the case when P is contained in $[0, 1]^n$, and a similar proof can be given in the more general case.

Proposition 2 (Properties 1, 2, 4) *Let M be admissible. Then $\partial M \subseteq GC \circ M$ and $\partial M \subseteq N_0$.*

Proof Given $T \subseteq \mathbb{R}^n$ a compact convex set, recall that $GC(T) = \bigcap_{\pi \in \mathbb{Z}^n} \{\pi x \leq \lfloor \sigma_T(\pi) \rfloor\}$. Let $P \subseteq \mathbb{R}^n$ be a polytope. First let $cx < d + 1$ with $c \in \mathbb{Z}^n$ and $d \in \mathbb{Z}$ be valid for $M(P)$. Then $cx \leq d$ is valid for $GC(M(P))$. It suffices to consider inequalities of this type even if $M(P)$ is not polyhedral by Lemma 2. As $M(P) \subseteq \{cx < d + 1\}$ it follows that $\emptyset = M(P) \cap \{cx \geq d + 1\} \supseteq M(P \cap \{cx \geq d + 1\})$. It follows that $cx \leq d$ is valid for $\partial M(P)$ and thus $\partial M(P) \subseteq GC(M(P))$.

Now let $P \subseteq [0, 1]^n$. For proving $\partial M(P) \subseteq N_0(P)$, recall that $N_0 = \bigcap_{i \in [n]} P_i$ where $P_i := \text{conv}\{(P \cap \{x_i = 0\}) \cup (P \cap \{x_i = 1\})\}$. We will show that $\partial M(P) \subseteq P_i$ for all $i \in [n]$. Therefore let $cx \leq d$ with $c \in \mathbb{Z}^n$ and $d \in \mathbb{Z}$ be valid for P_i with $i \in [n]$ arbitrary. (It is sufficient to consider only inequalities with integer coefficients since P_i is a rational polytope.) In particular, $cx \leq d$ is valid for $P \cap \{x_i = l\}$ with $l \in \{0, 1\}$. Thus we can conclude that $P \cap \{cx \geq d + 1\} \cap \{x_i = l\} = \emptyset$ for $i \in \{0, 1\}$. Therefore $x_i > 0$ and $x_i < 1$ are valid for $P \cap \{cx \geq d + 1\}$ and so by coordinate rounding (property (4.) of Definition 3), $x_i \leq 0$ and $x_i \geq 1$ are valid $M(P \cap \{cx \geq d + 1\})$. We obtain $M(P \cap \{cx \geq d + 1\}) = \emptyset$ and thus $cx \leq d$ is valid for $\partial M(P)$. \square

Remark 5 Note that, although the identity map I is not admissible as it does not satisfy property 4, it follows from the proof of Proposition 2 that $\partial I \subseteq GC$. Moreover in this particular case, it is not difficult to verify that $\partial I = GC$.

On the other hand, we remark that it is not true that the \mathbb{V} -scheme of an admissible operator M is always the composition of GC with M . For example, as shown in Sect. 5, $\partial GC(P) = P_I$ for any rational polytope in \mathbb{R}^2 . In contrast, it is well-known that $GC \circ GC(P) = GC^2(P)$ does not yield P_I for rational polytopes $P \subseteq \mathbb{R}^2$ in general. Also, as we will see later, the \mathbb{V} -schemes derive clique inequalities in a single round. This is not necessarily true for $GC \circ M(P)$ in general.

3.2 Comparing M and ∂M for M being GC, SC, N_0 , N , or N_+

We now compare various closures and their associated \mathbb{V} -closures. The first result shows that the verification scheme of the Gomory–Chvátal procedure is strictly stronger than split cuts.

Proposition 3 $\partial\text{GC} \subseteq \text{SC}$.

Proof We first verify that $\partial\text{GC} \subseteq \text{SC}$. Consider $cx \leq d$ being valid for $P \cap \{\pi x \leq \pi_0\}$ and $P \cap \{\pi x \geq \pi_0 + 1\}$ with $c, \pi \in \mathbb{Z}^n$ and $d, \pi_0 \in \mathbb{Z}$. Clearly, $cx \leq d$ is valid for $\text{SC}(P)$ and it suffices to consider inequalities $cx \leq d$ with this property as $\text{SC}(P)$ is a rational polytope [6] (since we work with polytopes, this is also implied by Lemma 2). Therefore consider $P \cap \{cx \geq d + 1\}$. By $cx \leq d$ being valid for the disjunction $\pi x \leq \pi_0$ and $\pi x \geq \pi_0 + 1$ we obtain that $P \cap \{cx \geq d + 1\} \cap \{\pi x \leq \pi_0\} = \emptyset$ and $P \cap \{cx \geq d + 1\} \cap \{\pi x \geq \pi_0 + 1\} = \emptyset$. This implies that $P \cap \{cx \geq d + 1\} \subseteq \{\pi x > \pi_0\}$ and similarly $P \cap \{cx \geq d + 1\} \subseteq \{\pi x < \pi_0 + 1\}$. We thus obtain that $\pi x \geq \pi_0 + 1$ and $\pi x \leq \pi_0$ are valid for $\text{GC}(P \cap \{cx \geq d + 1\})$. It follows $\text{GC}(P \cap \{cx \geq d + 1\}) = \emptyset$. Thus $cx \leq d$ is valid for $\partial\text{GC}(P)$.

To see that $\partial\text{GC} \subsetneq \text{SC}$, observe that $\partial\text{GC}(A_2) = \emptyset$ and $\text{SC}(A_2) \neq \emptyset$. □

Next we compare \mathbb{V} -schemes of two closures that are comparable. Before we present these results, we clarify the difference between the notion of *verifiable inequalities* against the notion of *valid inequalities for \mathbb{V} -closure of M* . Recall that given a polytope, $P \subseteq \mathbb{R}^n$, we say $cx \leq d$ is a verifiable inequality if $c \in \mathbb{Z}^n, d \in \mathbb{Z}$ and $M(P \cap \{cx \geq d + 1\}) = \emptyset$. Thus the \mathbb{V} -closure of M is the intersection of all verifiable inequalities. On the other hand, there may be a valid inequality for $\partial M(P)$ that is not verifiable. A trivial example of such as a valid inequality $cx \leq d$ for $\partial M(P)$ is when c is not a rational vector. The following example illustrates this difference more explicitly.

Example 2 Consider the set $P = \{x \in [0, 1]^2 \mid x_1 + x_2 \geq \frac{1}{2}, x_1 - x_2 \leq \frac{1}{2}, -x_1 + x_2 \leq \frac{1}{2}\}$. Observe that $\partial N_0(P) = P_I = \{(1, 1)\}$. This can be obtained by observing that the inequalities $x_1 \geq 1$ and $x_2 \geq 1$ are verifiable using N_0 . Now consider the inequality $2x_1 + 3x_2 \geq 5$. Clearly $2x_1 + 3x_2 \geq 5$ is valid for $\partial N_0(P)$ but is not verifiable since $N_0(P \cap \{2x_1 + 3x_2 \leq 4\}) \supseteq N_0(A_2) = \frac{1}{2}e$. □

The next result shows that switching to the verification schemes preserves inclusion. It holds for verification schemes of any two operators, as it is based on a purely geometric property.

Proposition 4 Let L, M be cutting-plane (not necessarily admissible) operators such that $L \subseteq M$. Then $\partial L \subseteq \partial M$.

Proof Let $P \subseteq \mathbb{R}^n$ be a polytope. By the definition of ∂M , it is sufficient to show that every inequality $cx \leq d$ verifiable by using M is valid for $\partial L(P)$. Now observe that since $cx \leq d$ is verifiable by using M , we have that $M(P \cap \{cx \geq d + 1\}) = \emptyset$. Thus, $L(P \cap \{cx \geq d + 1\}) = \emptyset$ since $L \subseteq M$ and therefore $cx \leq d$ is verifiable using L . Equivalently $cx \leq d$ is valid for $\partial L(P)$, completing the proof. □

In order to prove strict refinement or incompatibility between \mathbb{V} -closures the following proposition is helpful. It establishes when strict refinement carries over to the \mathbb{V} -schemes.

Proposition 5 (Properties 1, 3, 6) Let L, M be admissible. If $P \subseteq [0, 1]^n$ is a polytope with $P_I = \emptyset$ such that $M(P) = \emptyset$ and $L(P) \neq \emptyset$, then ∂L does not refine ∂M .

Before presenting the proof of Proposition 5, we first present a lemma which relates the \mathbb{V} -scheme with the actual closure.

Lemma 3 (Properties 1, 3, 6) *Let L be admissible. Let $P \subseteq [0, 1]^n$ be a polytope, $P \neq \emptyset$ and $P_I = \emptyset$. Define $Q \subseteq [0, 1]^{n+1}$ as $Q = \text{conv}(\{(x, 1) \in \mathbb{R}^{n+1} \mid x \in P\} \cup \{(y, 0) \in \mathbb{R}^{n+1} \mid y \in [0, 1]^n\})$. Then $\partial L(Q) = Q_I$ iff $L(P) = \emptyset$.*

Proof (\Leftarrow) If $L(P) = \emptyset$, then observe that $L(Q \cap \{x \in \mathbb{R}^{n+1} \mid x_{n+1} \geq 1\}) \cong L(P) = \emptyset$. Therefore $x_{n+1} \leq 0$ is valid for $\partial L(Q)$. Thus $\partial L(Q) = Q_I$.

(\Rightarrow) We will now show that if $L(P) \neq \emptyset$, then the point $\frac{1}{2}e \in \mathbb{R}^{n+1}$ satisfies $\frac{1}{2}e \in \partial L(Q)$ and hence $\partial L(Q) \neq Q_I$. Let $c \in \mathbb{Z}^n$ and $c_{n+1}, d \in \mathbb{Z}$ such that $L(T) = \emptyset$ with $T = \{x \in Q \mid cx + c_{n+1}x_{n+1} \geq d + 1\}$, that is $cx + c_{n+1}x_{n+1} \leq d$ is valid for $\partial L(Q)$. We will show that $c(\frac{1}{2}e^n) + \frac{1}{2}c_{n+1} \leq d$. Let $z_{min} := \min_{x \in [0, 1]^n} cx$ and $z_{max} := \max_{x \in [0, 1]^n} cx$; let x_{min} and x_{max} be a minimizer and a maximizer, respectively. Further we define $z_P := \min_{x \in P} cx$. As $L(P) \neq \emptyset$, by property 6 we have $P \cong Q \cap \{x_{n+1} = 1\} \not\subseteq T$ and therefore $z_P + c_{n+1} < d + 1$ and hence $c_{n+1} \leq d - z_P$. On the other hand we have that $z_{min} \leq z_P$ since $P \subseteq [0, 1]^n$. As $Q \cap \{x_{n+1} = 0\}$ is integral we have $T \cap (Q \cap \{x_{n+1} = 0\}) = \emptyset$ and so $z_{max} \leq d$. Moreover, by definition of x_{min} and x_{max} we can assume that x_{min} and x_{max} are antipodal, that is $e^n = x_{max} + x_{min}$. So we conclude

$$\frac{1}{2}e = \frac{1}{2}((x_{min}, 0) + (x_{max}, 0) + (0, \dots, 0, 1)),$$

and therefore

$$c \left(\frac{1}{2}e^n \right) + \frac{1}{2}c_{n+1} \leq \frac{1}{2}(z_P + d + (d - z_P)) = d$$

which completes the proof. □

We will use the following notation in the remainder of this section. Let $G \subseteq [0, 1]^n$ be a closed convex set. For $l \in [0, 1]$, by $G_{x_{n+1}=l}$ we denote the set $S \subseteq [0, 1]^{n+1}$ such that $S \cap \{x_{n+1} = l\} \cong G$ and S does not contain any other points. We can think of S arising from G by padding the coordinates of the points with l to the right. If G is the singleton $\{p\}$, then we write $\{p\}_{x_{n+1}=l}$ as $p_{x_{n+1}=l}$.

Proof of Proposition 5 Consider the auxiliary polytope Q given as $Q := \text{conv} \left(p_{x_{n+1}=1} \cup [0, 1]_{x_{n+1}=0}^n \right)$. By Lemma 3, $\partial L(Q) = \emptyset$ if and only if $L(P) \cong L(Q \cap \{x_{n+1} \geq 1\}) = \emptyset$ (and similarly for M). Since we have $M(P) = \emptyset$ but $L(P) \neq \emptyset$, we obtain $Q_I = \partial M(Q) \not\subseteq \partial L(Q)$. □

In the following propositions, polytopes are presented that help establish the strict inclusion or incompatibility depicted in Fig. 1, via Proposition 5.

Proposition 6 $\partial N_0 \perp \partial GC$ via the two polytopes $P_1 := \text{conv}([0, 1]^3 \cap \{x_1 + x_2 + x_3 = 3/2\}) \subseteq [0, 1]^3$ and $P_2 := \text{conv}(\{(\frac{1}{4}, \frac{1}{4}, 0), (\frac{1}{4}, \frac{1}{4}, 1), (\frac{1}{2}, 0, \frac{1}{2}), (\frac{1}{2}, 1, \frac{1}{2}), (0, \frac{1}{2}, \frac{1}{2}), (1, \frac{1}{2}, \frac{1}{2})\}) \subseteq [0, 1]^3$.

Proof By Proposition 5 it suffices to show that $GC(P_1) = \emptyset \neq N_0(P_1)$ and, vice versa, $GC(P_2) \neq \emptyset = N_0(P_2)$.

For the first case, clearly $GC(P_1) = \emptyset$. For proving that $N_0(P_1) \neq \emptyset$ it suffices to show that $\frac{1}{2}e$ is contained in $\text{conv}((P_1 \cap \{x_i = 0\}) \cup (P_1 \cap \{x_i = 1\}))$ for all $i \in [3]$. By symmetry, it suffices to show this for $i = 1$. This is true as $\frac{1}{2}e$ is the convex combination of the points $(0, 1, 1/2)$ and $(1, 0, 1/2)$.

For the second case, we first show that $N_0(P_2) = \emptyset$. For this observe that $\text{conv}((P_2 \cap \{x_3 = 0\}) \cup (P_2 \cap \{x_3 = 1\}))$ contains only points whose first two coordinates are equal to $1/4$. On the other hand

$$\begin{aligned} & \text{conv}((P_2 \cap \{x_1 = 0\}) \cup (P_2 \cap \{x_1 = 1\})) \\ & \cap \text{conv}((P_2 \cap \{x_2 = 0\}) \cup (P_2 \cap \{x_2 = 1\})) = \frac{1}{2}e, \end{aligned}$$

as $P_2 \cap \{x_3 = 1/2\} \cong A_2$ and thus $N_0(P_2) = \emptyset$. It thus remains to show that $GC(P_2) \neq \emptyset$. We will show that $\frac{1}{2}e \in P_2$. Let $cx \leq d$ with $c \in \mathbb{Z}^3$ be valid for P_2 . We divide the proof into two cases:

1. Either c_1 or c_2 is non-zero. In this case observe that

$$\begin{aligned} d & \geq d_0 := \max \left\{ c \left(\frac{1}{2}, 0, \frac{1}{2} \right), c \left(\frac{1}{2}, 1, \frac{1}{2} \right), c \left(0, \frac{1}{2}, \frac{1}{2} \right), c \left(1, \frac{1}{2}, \frac{1}{2} \right) \right\} \\ & > d_1 := c \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right), \end{aligned}$$

where the second inequality follows from the fact that $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ lies in the relative interior of the convex hull of $(\frac{1}{2}, 0, \frac{1}{2}), (\frac{1}{2}, 1, \frac{1}{2}), (0, \frac{1}{2}, \frac{1}{2}), (1, \frac{1}{2}, \frac{1}{2})$. Now observe that since $d_0, d_1 \in \frac{1}{2}\mathbb{Z}$, we obtain that the interval $[d_1, d_0]$ contains at least one integer number. Thus, $\lfloor d \rfloor \geq \lfloor d_0 \rfloor \geq d_1 = c(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$.

2. $c_1 = c_2 = 0$. If $c_3 > 0$, then $d \geq c_3$ (since $(\frac{1}{4}, \frac{1}{4}, 1) \in P_2$) and we obtain the GC inequality $c_3x_3 \leq \lfloor c_3 \rfloor$ where $\lfloor c_3 \rfloor \geq 1$. Thus this inequality cannot separate $\frac{1}{2}e$. Similarly if $c_3 \leq -1$, it can be verified that the resulting inequality cannot separate $\frac{1}{2}e$. □

Proposition 7 $\partial N_0 \perp \text{SC}$ via $P_1 := A_3 \subseteq [0, 1]^3$ and $P_2 := \text{conv}([0, 1]^3 \cap \{x_1 + x_2 + x_3 = 3/2\})$.

Proof Clearly $\text{SC} \not\subseteq \partial N_0$ as $\partial N_0(P_1) = \emptyset$ (proof similar to Example 1) but $\text{SC}(P_1) \neq \emptyset$ (cf. Lemma 3.3 in [8]).

For the converse, by Proposition 6 we have $N_0(P_2) \neq \emptyset$. However, $\text{SC}(Q) = Q_I$ by observing that the split $x_1 + x_2 + x_3 \leq 1$ and $x_1 + x_2 + x_3 \geq 2$ derives Q_I . Now the result follows from Proposition 5. □

Proposition 8 $\partial N \subsetneq \partial N_0$.

Proof We will show that there exists a polytope Q contained in the 0/1 cube such that $Q_I = \emptyset$ and $\emptyset = N(Q) \subsetneq N_0(Q)$. Then the result follows by the use of Proposition 5.

Let $P \subseteq [0, 1]^n$ such that $N(P) \subsetneq N_0(P)$, for example as discussed in page 171 of [16], for some $n \in \mathbb{N}$. Let $p \in N_0(P) \setminus N(P)$ and define

$$Q := \text{conv} \left(P_{x_{n+1}=1/2} \cup \{p_{x_{n+1}=1}, p_{x_{n+1}=0}\} \right).$$

Clearly, $Q_I = \emptyset$.

We first verify that $N(Q) = \emptyset$. Observe first that $N(Q) \subseteq N_0(Q) \subseteq \text{conv} (Q \cap \{x_{n+1} = 0\} \cup Q \cap \{x_{n+1} = 1\}) = \text{conv}(p_{x_{n+1}=0}, p_{x_{n+1}=1})$. On the other hand, it is easily verified that if $\sum_{i=1}^n c_i x_i \leq d$ is a valid inequality for P , then it is also a valid inequality for Q . Therefore we obtain that $N(Q) \subseteq \text{conv}(N(P)_{x_{n+1}=0}, N(P)_{x_{n+1}=1})$. Now since $\text{conv}(N(P)_{x_{n+1}=0}, N(P)_{x_{n+1}=1}) \cap \text{conv}(p_{x_{n+1}=0}, p_{x_{n+1}=1}) = \emptyset$, we obtain that $N(Q) = \emptyset$.

Next we verify that $N_0(Q) \neq \emptyset$. As $p \in N_0(P)$ we can conclude that $p_{x_{n+1}=1/2} \in \bigcap_{i \in [n]} \text{conv} (Q \cap \{x_i = 0\} \cup Q \cap \{x_i = 1\})$. Thus we have to show that $p_{x_{n+1}=1/2} \in \text{conv} (Q \cap \{x_{n+1} = 0\} \cup Q \cap \{x_{n+1} = 1\})$. This is clear though as $\{p_{x_{n+1}=1}, p_{x_{n+1}=0}\} \subseteq Q$. □

4 Rank of valid inequalities with respect to \mathbb{V} -closures

In this section, we establish several bounds on the rank of ∂M for the case of polytopes $P \subseteq [0, 1]^n$. Given a natural number k , we use the notation $M^k(P)$ and $\text{rk}_M(P)$ to be denote that k th closure of P with respect to M and the rank of P with respect to M respectively. As $\partial M \subseteq N_0$ we obtain:

Remark 6 [Upper bound in $[0, 1]^n$] (Properties 1, 2, 4) Let M be admissible and $P \subseteq [0, 1]^n$ be a polytope. Then $\text{rk}_{\partial M}(P) \leq n$.

Note that in general the property of M being admissible, does not guarantee that the upper bound on rank is n . For example, the GC closure can have a rank strictly higher than n (cf. [13,20]).

4.1 Rank of A_n

In quest for lower bounds on the rank of 0/1 polytopes, we note that among polytopes $P \subseteq [0, 1]^n$ that have $P_I = \emptyset$, the polytope $A_n = \{x \in [0, 1]^n \mid \sum_{i \in I} x_i + \sum_{i \notin I} (1 - x_i) \geq \frac{1}{2} \forall I \subseteq [n]\}$ has maximal rank (of n) for many admissible systems [18]. We will now establish that ∂M is not unrealistically strong by showing that it is subject to similar limitations. Recall that we do not prove *short verification* (property (7.)) for ∂M which is the basis for the lower bound in [19, Corollary 23] for admissible systems. We will show that the lower bound for ∂M is *inherited* from the original operator M . Let

$$F_n^k := \{x \in \{0, 1/2, 1\}^n \mid \text{exactly } k \text{ entries equal to } 1/2\},$$

and let $A_n^k := \text{conv} (F_n^k)$ be the convex hull of F_n^k . (Note $A_n^1 = A_n$.) With F being a face of $[0, 1]^n$ let $I(F)$ denote the index set of those coordinate that are fixed by F . We begin with a crucial lemma.

Lemma 4 (Properties 1, 2, 5, 6) *Let M be admissible and let $\ell \in \mathbb{N}$ such that $A_n^{k+\ell} \subseteq M(A_n^k)$ for all $n, k \in \mathbb{N}$ with $k + \ell \leq n$. If $n \geq k + 2\ell + 1$, then $A_n^{k+2\ell+1} \subseteq \partial M(A_n^k)$.*

Proof Let $P := A_n^k$ and let $cx \leq d$ with $c \in \mathbb{Z}^n$ and $d \in \mathbb{Z}$ be verifiable for $\partial M(P)$, that is $M(P \cap \{cx \geq d + 1\}) = \emptyset$. To prove this result, it is sufficient to prove that $A_n^{k+2\ell+1} \subseteq P \cap \{cx \leq d\}$.

We first claim that

$$A_{k+\ell}^k \cong A_n^k \cap F \not\subseteq P \cap \{cx \geq d + 1\} \tag{7}$$

for all $(k + \ell)$ -dimensional faces F of $[0, 1]^n$. Assume by contradiction that $A_n^k \cap F \subseteq P \cap \{cx \geq d + 1\}$. As $A_{k+\ell}^{k+\ell} \subseteq M(A_{k+\ell}^k)$ by assumption, we obtain $\emptyset \neq A_{k+\ell}^{k+\ell} \subseteq M(A_{k+\ell}^k) \cong M(A_n^k \cap F) \subseteq M(P \cap \{cx \geq d + 1\})$ which contradicts the verifiability of $cx \leq d$ over $\partial M(P)$.

Without loss of generality we can further assume that $c \geq 0$ and $c_i \geq c_j$ whenever $i \leq j$ by applying coordinate flips and permutations.

Next we claim that for all $(k + \ell)$ -dimensional faces F of $[0, 1]^n$, the point v^F defined as

$$v_i^F := \begin{cases} \in \{0, 1\} \text{ according to } F, & \text{for all } i \in I(F) \\ 0, & \text{if } c_i \text{ is one of the } \ell \text{ largest coefficients of } c \text{ with } i \notin I(F) \\ 1/2, & \text{otherwise} \end{cases} \tag{8}$$

for $i \in [n]$ is not contained in $P \cap \{cx \geq d + 1\}$, that is $cv^F < d + 1$ and so $cv^F \leq d + 1/2$. Note that $v^F \in P$ and observe that $v^F := \operatorname{argmin}_{x \in F_n^k \cap F} cx$. Therefore, if $v^F \in P \cap \{cx \geq d + 1\}$, then $A_n^k \cap F \subseteq P \cap \{cx \geq d + 1\}$ which in turn contradicts (7). This claim holds in particular for those faces F fixing coordinates to 1.

Finally, we claim that $A_n^{k+2\ell+1} \subseteq P \cap \{cx \leq d\}$. It suffices to show that $cv \leq d$ for all $v \in F_n^{k+2\ell+1}$ and we can confine ourselves to the worst case v given by

$$v_i := \begin{cases} 1, & \text{if } i \in [n - (k + 2\ell + 1)] \\ 1/2, & \text{otherwise.} \end{cases}$$

Observe that $cv \geq cw$ holds for all $w \in F_n^{k+2\ell+1}$. Let F be the $(k + \ell)$ -dimensional face of $[0, 1]^n$ obtained by fixing the first $n - (k + \ell)$ coordinates to 1. Then

$$\begin{aligned} cv &= \sum_{i=1}^{n-(k+2\ell+1)} c_i + \frac{1}{2} \sum_{i=n-(k+2\ell+1)+1}^n c_i \\ &\leq \sum_{i=1}^{n-(k+\ell)} c_i - \frac{1}{2}c_{n-(k+\ell)} + \sum_{i=n-(k+\ell)+1}^{n-k} 0 + \frac{1}{2} \sum_{i=(n-k)+1}^n c_i \\ &= cv^F - \frac{1}{2}c_{n-(k+\ell)} \leq d + \frac{1}{2} - \frac{1}{2}c_{n-(k+\ell)}. \end{aligned}$$

In case $c_{n-(k+\ell)} \geq 1$ it follows that $cv \leq d$. Therefore consider the case $c_{n-(k+\ell)} = 0$. Then we have that $c_i = 0$ for all $i \geq n - (k + \ell)$. In this case cv^F is integral and $cv^F < d + 1$ implies $cv^F \leq d$. So $cv \leq cv^F \leq d$ follows, which completes the proof. \square

Using Lemma 4 we can establish the following lower bound on the rank of ∂M for A_n .

Theorem 4 (Lower bound for A_n) (Properties 1, 2, 5, 6) *Let M be admissible and let $\ell \in \mathbb{N}$ such that $A_n^{k+\ell} \subseteq M(A_n^k)$ for all $n, k \in \mathbb{N}$ with $k + \ell \leq n$. If $n \geq k + 2\ell + 1$, then $rk_{\partial M}(A_n) \geq \lfloor \frac{n-1}{2\ell+1} \rfloor$.*

Proof We will show the $A_n^{1+k(2\ell+1)} \subseteq (\partial M)^k(A_n)$ as long as $n \geq k + 2\ell + 1$. The proof is by induction on k . Let $k = 1$, then $A_n^{1+2\ell+1} \subseteq \partial M(A_n^1) = \partial M(A_n)$ by Lemma 4. Therefore consider $k > 1$. Now $(\partial M)^k(A_n) = \partial M((\partial M)^{k-1}(A_n)) \supseteq \partial M(A_n^{1+(k-1)(2\ell+1)}) \supseteq A_n^{1+k(2\ell+1)}$, where the first inclusion follows by induction and the second inclusion by Lemma 4 again. Thus $(\partial M)^k(A_n) \neq \emptyset$ as long as $1 + k(2\ell + 1) \leq n$, which is the case as long as $k \leq \lfloor \frac{n-1}{2\ell+1} \rfloor$ and we can conclude $rk_{\partial M}(A_n) \geq \lfloor \frac{n-1}{2\ell+1} \rfloor$. \square

For $M \in \{GC, SC, N_0, N, N_+\}$ we have that $\ell = 1$ (see [19]) and therefore we obtain the following corollary.

Corollary 1 *Let $M \in \{GC, N_0, N, N_+, SC\}$ and $n \in \mathbb{N}$ with $n \geq 4$. Then $rk_{\partial M}(A_n) \geq \lfloor \frac{n-1}{3} \rfloor$.*

We can also derive an upper bound on the rank of A_n as follows.

Proposition 9 (Upper bound for A_n) (Properties 1, 2, 3, 4, 6) *Let M be admissible and $n \in \mathbb{N}$. Then $rk_{\partial M}(A_n) \leq n - 2$.*

Proof For $n \leq 3$, observe that the arguments presented in Example 1 for the case of ∂SC would be valid for any admissible cutting plane operator. Thus, the result holds for $n \leq 3$.

For $n \geq 4$, the proof is by induction on n . Consider $A_n \cap \{x_i = l\} \cong A_{n-1}$ for $(i, l) \in [n] \times \{0, 1\}$. Then after $n - 3$ applications of ∂M , by induction we have $(\partial M)^{(n-3)}(A_n \cap \{x_i = l\}) = \emptyset$. As $(i, l) \in [n] \times \{0, 1\}$ was arbitrary we obtain that $x_i < 1$ and $x_i > 0$ are valid for $(\partial M)^{(n-3)}(A_n)$. Another application of ∂M suffices to derive $x_i \leq 0$ and $x_i \geq 1$ and thus $(\partial M)^{(n-2)}(A_n) = \emptyset$ follows. \square

5 ∇ -closures for well-known and structured problems

We first establish a useful lemma which holds for any ∂M with M being admissible. The lemma is analogous to Lemma 1.5 in [16].

Lemma 5 (Properties 1, 2, 4) *Let M be admissible, let $P \subseteq [0, 1]^n$ be a closed convex set and let $(c, d) \in \mathbb{Z}_+^{n+1}$. If $cx \leq d$ is valid for $P \cap \{x_i = 1\}$ for every $i \in [n]$ with $c_i > 0$, then $cx \leq d$ is valid for $\partial M(P)$.*

Proof Clearly, $cx \leq d$ is valid for P_I : if $x \in P \cap \mathbb{Z}^n$ is non-zero, then there exists an $i \in [n]$ with $x_i = 1$, otherwise $cx \leq d$ is trivially satisfied.

We claim that $cx \leq d$ is valid for $\partial M(P)$. Let $Q := P \cap \{cx \geq d + 1\}$ and observe that $Q \cap \{x_i = 1\} = \emptyset$ for any $i \in [n]$ with $c_i > 0$. Therefore by the coordinate rounding property of admissible operators, we have that $M(Q) \subseteq \bigcap_{i \in [n]; c_i > 0} \{x_i = 0\}$. By definition of Q we also have that $M(Q) \subseteq \{cx \geq d + 1\}$. Since $c \geq 0$ and $d \geq 0$ we deduce $M(Q) = \emptyset$ and the claim follows. \square

5.1 Monotone polytopes

The following theorem is a direct consequence of Lemma 5 and follows in a similar fashion as Lemma 2.7 in [5] or Lemma 2.14 in [16].

Theorem 5 (Properties 1, 2, 3, 4) *Let M be admissible. Further, let $P \subseteq [0, 1]^n$ be a polytope and $(c, d) \in \mathbb{Z}_+^{n+1}$ such that $cx \leq d$ is valid for $P \cap F$ whenever F is an $(n - k)$ -dimensional face of $[0, 1]^n$ obtained by fixing coordinates to 1. Then $cx \leq d$ is valid $(\partial M)^k(P)$.*

Proof The proof is by induction on k , the number of coordinates fixed to obtain a $n - k$ dimensional face. For $k = 1$ the assertion follows with Lemma 5. Therefore let $k > 1$. Define $Q_i = P \cap \{x_i = 1\}$ for all $i \in [n]$. Then $cx \leq d$ is valid for $Q_i \cap \tilde{F}$ whenever \tilde{F} is an $(n - 1) - (k - 1)$ -dimensional face of $[0, 1]^{n-1}$ fixing $k - 1$ coordinates to 1 and i is not one of those coordinates. We can apply the induction hypothesis obtaining that $cx \leq d$ is valid for $(\partial M)^{k-1}(Q_i)$ for all $i \in [n]$. By homogeneity of ∂M we obtain $(\partial M)^{k-1}(Q_i) = (\partial M)^{k-1}(P) \cap \{x_i = 1\}$ for all $i \in [n]$. Applying Lemma 5 once more yields that $cx \leq d$ is valid for $(\partial M)^k(P)$. \square

We call a polytope $P \subseteq [0, 1]^n$ *monotone* if $x \in P$, $y \in [0, 1]^n$, and $y \leq x$ (coordinate-wise) implies $y \in P$. We can derive the following corollary from Theorem 5 which is the analog to Lemma 2.7 in [5].

Corollary 2 (Properties 1, 2, 3, 4) *Let M be admissible and let $P \subseteq [0, 1]^n$ be a monotone polytope with $\max_{x \in P_I} ex = k$. Then $rk_{\partial M}(P) \leq k + 1$.*

Proof Observe that since P is monotone, so is P_I and thus P_I possesses an inequality description $P = \{x \in [0, 1]^n \mid Ax \leq b\}$ with $A \in \mathbb{Z}_+^{m \times n}$ and $b \in \mathbb{Z}_+^m$ for some $m \in \mathbb{N}$. Therefore it suffices to consider inequalities $cx \leq d$ valid for P_I with $c, d \geq 0$. As $\max_{x \in P_I} ex = k$ and P is monotone, we claim that $P \cap F = \emptyset$ whenever F is an $n - (k + 1)$ dimensional face of $[0, 1]^n$ obtained by fixing $k + 1$ coordinates to 1. Assume by contradiction that $x \in P \cap F \neq \emptyset$. As $P \cap F$ is monotone, the point obtained by setting all fractional entries of x to 0 is contained in $P_I \cap F$ which is a contradiction to $\max_{x \in P_I} ex = k$. Therefore $cx \leq d$ is valid for all $P \cap F$ with F being an $n - (k + 1)$ dimensional face of $[0, 1]^n$ obtained by fixing $k + 1$ coordinates to 1. The result follows now by using Theorem 5. \square

5.2 Stable set polytope

Given a graph $G := (V, E)$, the fractional stable set polytope of G is given by $FSTAB(G) := \{x \in [0, 1]^n \mid x_u + x_v \leq 1 \forall (u, v) \in E\}$. See [16] for a description of clique inequalities, odd hole inequalities, odd anti-hole inequalities, and odd wheel inequalities for the stable set polytope.

Now Lemma 5 can be used to prove the following result.

Theorem 6 (Properties 1, 2, 4) *Clique inequalities, odd hole inequalities, odd anti-hole inequalities, and odd wheel inequalities are valid for $\partial M(FSTAB(G))$ with M being an admissible operator.*

Proof In the following let

$$P^0 := \left\{ x \in [0, 1]^{|V|} \mid x_u + x_v \leq 1 \forall (u, v) \in E \right\},$$

for V, E chosen as explained below.

We first consider the clique inequalities. Let $H(V, E)$ be an induced clique. Then the clique inequality is

$$\sum_{u \in V} x_u \leq 1.$$

Now for every vertex v in V , fixing $x_v = 1$ in the system P^0 implies that $x_u = 0$ for $u \neq v$. Thus, the clique inequality is valid for $P^0 \cap \{x_v = 1\}$ for all $v \in V$. Now by Lemma 5 the result follows.

Odd hole inequalities are GC inequalities: Add all the inequalities of the form $x_u + x_v \leq 1$ along the odd hole, divide by 2, and the round down the right-hand-side. Therefore, odd hole inequalities are valid for ∂M .

Let $H(V, E)$ be an induced graph which is an odd anti-hole with $|V| \geq 5$. Call the complement of $H(V, E)$ as $\bar{H}(V, \bar{E})$ (i.e. \bar{H} is an odd hole). Then the odd anti-hole inequality is

$$\sum_{u \in V} x_u \leq 2.$$

For every vertex v in V , fixing $x_v = 1$ in the system

$$P^0 = \left\{ x \in [0, 1]^{|V|} \mid x_u + x_v \leq 1 \forall (u, v) \in E \right\},$$

implies that $x_u = 0$ for all u except the neighbors of vertex v in \bar{H} . Moreover, these two neighbors of v in \bar{H} are neighbors of each other in H (since $|V| \geq 5$). Thus, $\max \sum_{u \in V} x_u = 2$ for $x \in P^0 \cap \{x_v = 1\}$. Now by Lemma 5 the result follows.

Let $H(\{0, \dots, n\}, E)$ be an induced graph which is an odd wheel, that is n is odd, the vertices 1 through n form a hole and the vertex 0 is a neighbor to all other vertices. Then the odd wheel inequality is

$$\sum_{i=1}^n x_i + \frac{n-1}{2}x_0 \leq \frac{n-1}{2}.$$

Now for the vertex 0, fixing $x_0 = 1$ in the system

$$P^0 = \left\{ x \in [0, 1]^{n+1} \mid x_u + x_v \leq 1 \forall (u, v) \in E \right\},$$

implies that $x_u = 0$ for $u \in \{1, \dots, n\}$. Therefore, $\max \sum_{i=1}^n x_i + \frac{n-1}{2}x_0 = \frac{n-1}{2}$ for $x \in P^0 \cap \{x_0 = 1\}$.

On fixing $x_1 = 1$ in P^0 , we obtain that $x_0 = 0, x_2 = 0, x_n = 0$ and therefore the system P^0 reduces to

$$x_k + x_{k+1} \leq 1 \forall k \in \{2, \dots, n-2\} \tag{9}$$

$$0 \leq x_k \leq 1 \forall k \in \{2, \dots, n-2\}. \tag{10}$$

Now observe that the constraint set (9) is totally unimodular. Therefore, $\max \sum_{i=1}^n x_i + \frac{n-1}{2}x_0 = \frac{n-1}{2}$ for $x \in P^0 \cap \{x_1 = 1\}$. Similarly, $\max \sum_{i=1}^n x_i + \frac{n-1}{2}x_0 = \frac{n-1}{2}$ for $x \in P^0 \cap \{x_v = 1\}$ for $v \in \{2, \dots, n\}$. Now by Lemma 5 the result follows. □

5.3 The traveling salesman problem

So far we have seen that transitioning from a general cutting-plane procedure M to its \mathbb{V} -scheme, ∂M , can result in a significantly lower rank for valid inequalities, potentially making them accessible in a small number of rounds. However, we will now show that the rank of (the subtour elimination relaxation of) the traveling salesman polytope remains high, even when using \mathbb{V} -schemes of strong operators such as SC or N_+ . For $n \in \mathbb{N}$, let $G = (V, E)$ be the complete graph on n vertices and $H_n \subseteq [0, 1]^n$ be the polytope given by (see [5] for more details)

$$\begin{aligned} x(\delta(\{v\})) &= 2 && \forall v \in V \\ x(E(W)) &\leq |W| - 1 \forall \emptyset \subsetneq W \subsetneq V \\ x_e &\in [0, 1] && \forall e \in E, \end{aligned}$$

where for a given node v , $x(\delta(\{v\}))$ is the sum of the components of the vector x corresponding to edges incident to the node v and for any subset W of V , $x(E(W))$ is the sum of the components of the vector x corresponding to edges which are incident to nodes contained only in W . Note that the dimension of H_n is $\Theta(n^2)$. We obtain the following statement which is the analog to [5, Theorem 4.1]. A similar result for the admissible systems M in general can be found in the full-length version of [19].

Theorem 7 (Properties 1, 2, 3, 4, 5, 6) *Let $M \in \{GC, N_0, N, N_+, SC\}$. For $n \in \mathbb{N}$ and H_n as defined above we have $rk_{\partial M}(H_n) \in \Theta(n)$. In particular $rk_{\partial M}(H_n) \in \Theta(\sqrt{\dim(P)})$.*

Proof We first establish the lower bound. As shown in [3] or [5, Theorem 4.1], there exists an embedding

$$f : A_{\lfloor n/8 \rfloor} \hookrightarrow H_n,$$

consisting of coordinate flips and coordinate duplications only, such that $f(\frac{1}{2}e) \in H_n \setminus (H_n)_I$. Since ∂M is almost admissible, we have that ∂M commutes with f . We obtain

$$f\left(\frac{1}{2}e\right) \in f(\partial M^k(A_{\lfloor n/8 \rfloor})) = \partial M^k(f(A_{\lfloor n/8 \rfloor})) \subseteq \partial M^k(H_n),$$

for $k < \text{rk}_{\partial M}(A_{\lfloor n/8 \rfloor})$ and thus $\text{rk}_{\partial M}(H_n) \geq \text{rk}_{\partial M}(A_{\lfloor n/8 \rfloor}) \in \Omega(n)$ by Corollary 1.

For the upper bound, observe that H_n is a face of T_n given by

$$\begin{aligned} x(\delta(\{v\})) &\leq 2 && \forall v \in V \\ x(E(W)) &\leq |W| - 1 && \forall \emptyset \subsetneq W \subsetneq V \\ x_e &\in [0, 1] && \forall e \in E. \end{aligned}$$

(see [5] for details). As T_n is given by a system of inequalities of the form $Ax \leq b$ with non-negative coefficients, it follows that T_n is a monotone polytope. Furthermore, we can conclude that $\max_{x \in (T_n)_I} ex \leq n$ so that we can apply Corollary 2. We obtain that $\text{rk}_{\partial M}(H_n) \leq \text{rk}_{\partial M}(T_n) \leq n + 1$ which finishes the proof. \square

The same result can be shown to hold for the asymmetric TSP problem (see [3] and [5]).

5.4 General polytopes in \mathbb{R}^2

The GC rank of valid inequalities for polytopes in \mathbb{R}^2 can be arbitrarily high; see example in [17]. The SC rank of valid inequalities for polytopes in \mathbb{R}^2 can be at least 2; A_2 is an example where the split rank is 2 and the instance is infeasible and see [12] for an example where the instance is feasible and the split rank is at least 2.

However, ∂GC is significantly stronger as shown next.

In the following proof, a *split* is a set of the form $\{g \leq hx \leq g + 1\}$ where $h \in \mathbb{Z}^2$ and $g \in \mathbb{Z}$ and we call the lines $\{hx = g\}$ and $\{hx = g + 1\}$ as the boundary lines of the split. A set $Q \subset \mathbb{R}^2$ is called *lattice-free* if $\text{int}(Q) \cap \mathbb{Z}^2 = \emptyset$. Therefore a split set is an example of a lattice-free convex set.

The following result follows from [15].

Lemma 6 *If $P \subseteq \mathbb{R}^2$ is a full-dimensional unbounded lattice-free convex set, then P is contained in a split set.*

Theorem 8 *Let P be a polytope in \mathbb{R}^2 . Then $\partial \text{GC}(P) = P_I$.*

Proof The proof is divided into various cases based on the dimension of P_I .

Case 1 $\dim(P_I) = 2$. We will show that every facet-defining inequality can be obtained using the ∂GC operator. Given a facet-defining inequality $cx \leq d$, there exists at least two integer points belonging to P_I that satisfy the inequality at equality. Let $Q := P \cap \{x \in \mathbb{R}^2 \mid cx \geq d\}$. We assume that $Q \not\subseteq \{cx < d + 1\}$, since otherwise $cx \leq d$ is a GC cut. Let x^1 and x^2 be two consecutive integer points on the face of P_I defined by $\{cx = d\}$. As $Q \not\subseteq \{cx < d + 1\}$ we obtain that Q intersects $\{cx = d + 1\}$ in a segment contained between two integer points y^1, y^2 on $\{cx = d + 1\}$. Since x^1, x^2, y^2, y^1 (in topological order) are the vertices of a lattice-free parallelogram and these are the only integer points contained in this parallelogram, the lines l^1 and l^2 through x^1y^1 and x^2y^2 respectively are the boundary lines of a split. Call this split S . It is sufficient to verify that $Q \cap \{cx \geq d + 1\}$ is strictly contained in S , since this implies that $\text{GC}(Q \cap \{cx \geq d + 1\}) = \emptyset$. Note that Q is contained in the union of the split set $d \leq cx \leq d + 1$ and S , since otherwise it is straightforward to verify that y^i for some $i \in \{1, 2\}$ must belong to Q . Therefore $Q \cap \{cx \geq d + 1\}$ is contained in S , and it remains to prove that $Q \cap \{cx \geq d + 1\}$ is strictly contained in S . Let $\hat{l}^i \subseteq l^i$ be the half-line starting at the point y^i which does not contain the point x^i . Since $x^i \in P$ and $y^i \notin P$, by convexity of P we have that $\hat{l}^i \cap P = \emptyset$. Moreover, since $Q \subseteq P$, we have that $Q \cap \{cx \geq d + 1\} \cap \hat{l}^i = \emptyset$ for $i \in \{1, 2\}$. Since $Q \cap \{cx \geq d + 1\}$ is contained in S , we therefore obtain that $Q \cap \{cx \geq d + 1\}$ is strictly contained in S . \diamond

Before we consider other cases where $\dim(P_I) \leq 1$, observe that since P is a rational polytope, with out loss of generality we may assume that P is full-dimensional: Since P is rational polytope, it is straightforward to verify that there exists a full-dimensional rational polytope T satisfying $P \subseteq T$ and $T_I = P_I$. Since $\partial\text{GC}(P) \subseteq \partial\text{GC}(T)$ it is sufficient to verify that $\partial\text{GC}(T) = T_I$.

If P contains integer points, then we further preprocess P in the following fashion. Suppose there exists a linear inequality in the description of P , such that removing this inequality from the description of P results in a polyhedron P' such that $P'_I = P_I$. Note that in this case P' is a polytope since P' is rational and $P'_I = P_I \neq \emptyset$. We rename P' by P and by applying this procedure iteratively, we obtain a polytope where removing any facet-defining inequality introduces new integer points.

We call a point $v \in \mathbb{Z}^2$ as *minimally infeasible for facet-defining inequality* $fx \leq g$ of P if (1) v satisfies all the constraints defining P except $fx \leq g$ and (2) $\text{conv}(P \cup \{v\}) \cap \mathbb{Z}^2 = (P \cap \mathbb{Z}^2) \cup \{v\}$. A minimally infeasible point exists for every facet-defining inequality of P : By the preprocessing of P , we know that removing the facet-defining inequality $fx \leq g$ introduces new integer points in P . Now it is straightforward to verify that any optimal solution of the problem $\min\{fx \mid x \in \hat{P} \cap \mathbb{Z}^2\}$ is a minimally infeasible point for $fx \leq g$, where \hat{P} is the set obtained by replacing $fx \leq g$ by $fx > g$ in the description of P .

Case 2 $\dim(P_I) = 1$. Let $P_I \subseteq \{cx = d\}$ where $c \in \mathbb{Z}^2$ and $d \in \mathbb{Z}$. Note that the inequality $cx \leq d$ satisfies at least two integer points belonging to P_I at equality. Therefore by the application of the proof technique used in case 1, we have that the inequality $cx \leq d$ is valid for $\partial\text{GC}(P)$. Similarly, $cx \geq d$ is valid for $\partial\text{GC}(P)$. Thus,

$$\partial\text{GC}(P) \subseteq \{cx = d\}. \tag{11}$$

Let $P_I = \{cx = d, g \leq fx \leq h\}$, where f is a vector orthogonal to c . Next observe that there exists at least one facet-defining inequality of P that separates the integer points in $\{cx = d, fx < g\}$ from P . Call one such facet-defining inequality as F^1 . Similarly let F^2 be a facet-defining inequality that separates the integer points in $\{cx = d, fx > h\}$ from P . Since P is full-dimensional, bounded and non-empty it has at least three facets. Select a facet-defining inequality of P different from F^1 and F^2 . Let $v \in \mathbb{Z}^2$ be a minimally infeasible point for this facet and let $P' := \text{conv}(P \cup \{v\})$. Since F^1 and F^2 still separate the integer points in the the set $\{cx = d, fx < g\} \cup \{cx = d, fx > h\}$ from P' , we may assume without loss of generality

$$v := (P' \setminus P) \cap \mathbb{Z}^2 \text{ satisfies } cx < d. \tag{12}$$

Therefore, $\dim(\text{conv}(P'_I)) = 2$ and by case 1, we have that $\partial\text{GC}(P') = P'_I$. Finally note that since $\partial\text{GC}(P) \subseteq \partial\text{GC}(P')$, we obtain that

$$\partial\text{GC}(P) \subseteq \{cx = d\} \cap \partial\text{GC}(P') \tag{13}$$

$$= \{cx = d\} \cap P'_I \tag{14}$$

$$= P_I, \tag{15}$$

where (13) follows from (11) and (15) follows from (12).

Case 3 $\dim(P_I) = 0$. Since P is full-dimensional, bounded and non-empty it has at least three facets. Let v^1 and v^2 be minimally infeasible points for two distinct facet-defining inequalities for P . Then we have that $v^1 \neq v^2$. Let $P^i = \text{conv}(P \cup \{v^i\})$ for $i \in \{1, 2\}$. Then we have $\partial\text{GC}(P) \subseteq \partial\text{GC}(P^1 \cap P^2) \subseteq \partial\text{GC}(P^1) \cap \partial\text{GC}(P^2) = P^1_I \cap P^2_I = \{u\}$, where the second last equality follows from case 2 and the fact that $\dim(P^i_I) = 1$ and the last equality follows from the fact that $v^2 \notin \text{conv}(\{v^1\} \cup \{u\})$ and $v^1 \notin \text{conv}(\{v^2\} \cup \{u\})$.

Case 4 $P_I = \emptyset$: Like in the case where $P_I \neq \emptyset$, we preprocess P by removing any facet-defining inequality if removing the inequality does not introduce any integer points in the resulting polyhedron. Note that, in this case, the resulting set need not be bounded. If P is unbounded, since it is lattice-free and full-dimensional, by Lemma 6 we have that P is contained in a split set $\{a_0 \leq ax \leq a_0 + 1\}$. Moreover since $P \cap \mathbb{Z}^2 = \emptyset$, we obtain that $P \cap \{ax \geq a_0 + 1\}$ is a line segment (possibly an empty set) strictly contained between two integer points. This implies that $\text{GC}(P \cap \{ax \geq a_0 + 1\}) = \emptyset$ and thus $ax \leq a_0$ is a valid inequality for $\partial\text{GC}(P)$. Similarly, $ax \geq a_0 + 1$ is a valid inequality for $\partial\text{GC}(P)$, completing the proof.

Now consider the case where P is bounded. Then P has at least three facets. Let v^1 and v^2 be minimally infeasible points for two distinct facet-defining inequalities of P and let $P^i := \text{conv}(\{v^i\} \cup P)$ for $i \in \{1, 2\}$. Therefore, $\partial\text{GC}(P) \subseteq \partial\text{GC}(P^1 \cap P^2) \subseteq \partial\text{GC}(P^1) \cap \partial\text{GC}(P^2) = \{v^1\} \cap \{v^2\} = \emptyset$ where the second last equality follows from case 3 and the last equality follows from the fact that $v^1 \neq v^2$. □

6 Concluding remarks

In this paper, we consider a new paradigm for generating cutting-planes. Rather than *computing* a cutting-plane we suppose that the cutting-plane is given, either by a *deliberate construction* or guessed in some other way and then we *verify* its validity using a regular cutting-plane procedure. We have shown that cutting-planes obtained via the verification scheme can be very strong, significantly exceeding the capabilities of the regular cutting-plane procedure. This superior strength is illustrated, for example, in Theorems 1, 3, Fig. 1, Lemma 6, Proposition 9, Theorems 5, 6, 7 and 8. On the other hand, we also show that the verification scheme is not unrealistically strong, as illustrated by Theorems 4 and 7.

We would like to point out that verification schemes (with minor adjustments) can also be applied to mixed-integer programming problems to generate pure integer cuts. For example one could replace the Gomory–Chvátal generator with projected Gomory–Chvátal cuts in the mixed-integer case.

Acknowledgments The authors are most grateful to the anonymous referees for their detailed remarks that considerably improved the presentation and simplified some of the proofs.

References

1. Balas, E., Ceria, S., Cornuéjols, G.: A lift-and-project cutting plane algorithm for mixed integer 0–1 programs. *Math. Program.* **58**, 295–324 (1993)
2. Chvátal, V.: Edmonds polytopes and a hierarchy of combinatorial problems. *Discret. Math.* **4**, 305–337 (1973)
3. Chvátal, V., Cook, W., Hartmann, M.: On cutting-plane proofs in combinatorial optimization. *Linear Algebra Appl.* **114**, 455–499 (1989)
4. Cook, W., Coullard, C.R., Turan, G.: On the complexity of cutting plane proof. *Math. Program.* **47**, 11–18 (1990)
5. Cook, W., Dash, S.: On the matrix cut rank of polyhedra. *Math. Oper. Res.* **26**, 19–30 (2001)
6. Cook, W., Kannan, R., Schrijver, A.: Chvátal closures for mixed integer programming problems. *Math. Program.* **58**, 155–174 (1990)
7. Cornuéjols, G., Li, Y.: Elementary closures for integer programs. *Oper. Res. Lett.* **28**, 1–8 (2001)
8. Cornuéjols, G., Li, Y.: On the rank of mixed 0–1 polyhedra. *Math. Program.* **91**, 391–397 (2002)
9. Dadush, D., Dey, S.S., Vielma, J.P.: On the Chvátal–Gomory closure of a compact convex set. In: Günlük, O., Woeginger, G.J. (eds.) IPCO 2011, Proceedings Lecture Notes in Computer Science, vol. 6655, pp. 130–142. Springer, Berlin (2011)
10. Dadush, D., Dey, S.S., Vielma, J.P.: The Chvátal–Gomory closure of strictly convex body. *Math. Oper. Res.* **36**, 227–239 (2011)
11. Dey, S.S., Pokutta, S.: Design and verify: a new scheme for generating cutting-planes. In: Günlük, O., Woeginger, G.J. (eds.) IPCO 2011, Proceedings Lecture Notes in Computer Science, vol. 6655, pp. 143–155. Springer, Berlin (2011)
12. Dey, S.S., Richard, J.P.P.: Some relations between facets of low- and high-dimensional group problems. *Math. Program.* **123**, 285–313 (2010)
13. Eisenbrand, F., Schulz, A.S.: Bounds on the Chvátal rank of polytopes in the 0/1-cube. *Combinatorica* **23**, 245–262 (2003)
14. Gomory, R.E.: Outline of an algorithm for integer solutions to linear programs. *Bull. Am Math. Soc.* **64**, 275–278 (1958)
15. Lovász, L.: Geometry of numbers and integer programming. *Mathematical Programming: Recent Developments and Applications* (1989)
16. Lovász, L., Schrijver, A.: Cones of matrices and set-functions and 0–1 optimization. *SIAM J. Optim.* **1**, 166–190 (1991)

17. Nemhauser, G.L., Wolsey, L.A.: *Integer and Combinatorial Optimization*. Wiley-Interscience, New York (1988)
18. Pokutta, S., Schulz, A.S.: Characterization of integer-free 0/1 polytopes with maximal rank (xxxx) (submitted)
19. Pokutta, S., Schulz, A.S.: On the rank of generic cutting-plane proof systems. In: Eisenbrand, F., Shepherd, B. (eds.) *Integer Programming and Combinatorial Optimization, 14th International IPCO Conference, Proceedings, Lecture Notes in Computer Science*, Lausanne, Switzerland, June 9–11, 2010, pp. 450–463. Springer, Berlin (2010)
20. Pokutta, S., Stauffer, G.: Lower bounds for the Chvátal-Gomory rank in the 0/1 cube. *Oper. Res. Lett* (2011) (to appear)
21. Schrijver, A.: *Theory of Linear and Integer Programming*. Wiley, New York (1986)

Copyright of Mathematical Programming is the property of Springer Science & Business Media B.V. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.