

BRIEF REPORT

Multimedia decision support intervention: A promising approach to enhance the intention to complete an advance directive among hospitalized adults

Ronald L. Hickman Jr., PhD, RN, ACNP-BC (Assistant Professor)^{1,2}, Amy R. Lipson, PhD (Research Associate)², Melissa D. Pinto, PhD, RN (Assistant Professor)³, & Grant Pignatiello, BSN, RN (Research Assistant)¹

¹School of Nursing, Case Western Reserve University, Cleveland, Ohio

²University Hospitals Case Medical Center, Cleveland, Ohio

³School of Nursing, Emory University, Atlanta, Georgia

Keywords

Advance directives; hospitalized adults; educational interventions; decision support; critical illness.

Correspondence

Ronald L. Hickman, Jr., PhD, RN, ACNP-BC, School of Nursing, Case Western Reserve University, 2120 Cornell Road, Cleveland, OH 44106-4904. Tel: 216-832-5296 (home), 216-368-2147 (work); Fax: 216-368-5989; E-mail: rlh4@case.edu

Received: February 2013;
accepted: May 2013

doi: 10.1002/2327-6924.12051

Abstract

Purpose: We examined the acceptability and initial efficacy of a multimedia decision support (MDS) intervention to improve intention to complete an advanced directive (AD) among hospitalized adults following an episode of critical illness.

Data sources: We used comparative quasi-experimental posttest only design. Forty-nine hospitalized adults, recovering from a critical illness, received either MDS or AD educational brochure. Demographic characteristics and self-report measures of AD knowledge were captured at baseline and used as covariates. Helpfulness of the intervention (acceptability) and the outcome variable, intention to complete an AD decision, were assessed after exposure to the MDS intervention or educational brochure (information-only control condition).

Conclusions: The MDS was a more acceptable form of education compared to a brochure. After adjusting for covariates, participants exposed to the MDS intervention were 24.7 times more likely to intend to complete an AD compared to those who were assigned to the information-only control condition.

Implications for practice: This pilot study establishes the acceptability and initial efficacy of the MDS intervention among individuals with critical illness, who are at high risk for hospital readmission life-sustaining treatment. This study illuminates a teachable moment in which patients are more receptive to interventions to complete an AD.

The Patient Protection and Affordable Healthcare and Federal Patient Self-Determination Acts underscore the importance of decision support and advance care planning. However, Kass-Bartelmes, Hughes, and Rutherford (2003) and Camhi et al. (2009) report that less than 50% of hospitalized adults who were severely or terminally ill had advance directives (ADs) documented in their medical records. Hospitalized adults who are critically ill and require temporary life-sustaining treatments, such as mechanical ventilation, hemodialysis, or artificial nutrition, represent a high-risk cohort likely to experience frequent hospital readmissions, impaired functional status, and a recurring need for life-sustaining treatments

(Cox & Carson, 2007; Cox, Carson, & Lindquist, 2007; Cox et al., 2009; Daly & Douglas, 2005; Douglas & Daly, 2001; Douglas, Daly, Gordon, & Brennan, 2002; Douglas, Daly, Kelley, O'Toole, & Montenegro, 2007). Among chronically critically ill adults, AD completion rates are low. These individuals are at high risk for readmission. In a recent study by Camhi et al. (2009), only 16% of chronically ill adults with prolonged mechanical ventilation had ADs established before or after their episode of critical illness. This research illuminates the need for establishing ADs prior to the patient's discharge and before subsequent hospital admissions to promote patient-centered decision making. Yet, few studies have explored

educational interventions to promote intention to complete an AD among hospitalized adults recovering from a critical illness.

A patient's perception of illness and their preferences for life-sustaining treatments may change as a result of their hospital experience. Exposure to life-sustaining treatments and convalescence in a hospital may serve as opportune time to provide AD education to hospitalized adults. Teachable moments, naturally occurring cueing events, can alter an individual's state of receptiveness towards receiving health education and stimulate their contemplation, leading to a desired health behavior change (Cohen, Clark, Lawson, Casucci, & Flocke, 2011; Flocke et al., 2012). Previous work among patients admitted to an emergency department or an intensive care unit (ICU) for an acute health problems (Boudreaux, Bock, & O'Hea, 2012; Boudreaux et al., 2010; Clark & Moss, 2011), and several studies conducted in primary care and outpatient settings indicate that the risk of acute health problems can signify a teachable moment that provides healthcare professionals an opportunity to introduce health education resources and effectively facilitate a desired health behavior change (Cohen et al., 2011; Floyd, Steffens, Pavlik, & Andrykowski, 2011; Sommers, Lyons, Bohn, Ribak, & Fargo, 2013). Building on this previous research and the premise of a teachable moment, we postulated that the experience of surviving an episode of critical illness requiring an acute episode of mechanical ventilation may alter a hospitalized adult's perspective on advance care planning by increasing their receptiveness to educational interventions focused on AD completion.

This pilot study aimed to capitalize on the routinely missed teachable moment by examining the effects of two educational interventions on established factors that influence AD decision making (age, race, and knowledge of ADs) and the intention to complete AD among hospitalized adults (Alano et al., 2010; Salmond & Estrellam, 2005; Tamura, Goldstien, & Perez-Stable, 2010). Therefore, the purpose of this study is to present evidence on the acceptability and initial efficacy of a multimedia decision support (MDS) intervention when compared to the American Hospital Association's commonly used educational brochure, *Putting It In Writing*, among hospitalized adults recovering from a critical illness.

Methods

Design

As shown in Figure 1, this is a comparative quasi-experimental study with a posttest-only design. The first 30 subjects enrolled into this study were assigned to the

control condition and received the American Hospital Association's *Putting It In Writing* brochure (retrieved from <http://www.aha.org/content/13/putitinwriting.pdf>). Subsequently enrolled subjects ($n = 19$) were assigned to the experimental condition, the MDS intervention. Subjects exposed to the MDS intervention received a single dose of the interactive computer program administered on a laptop computer. The components of the intervention were (a) a 10-min video vignette of a patient and his spouse who have not considered an AD and conveys the psychological impact of not having an AD had on the spouse while providing education about ADs, and (b) interactive education consisting of state specific AD education and additional internet resources on advance care planning. Subjects assigned to the either study condition were exposed a single of dose of the respective educational intervention immediately after the collection of their demographic data.

Sample

A convenience sample of hospitalized adults ($N = 49$) who required an episode of acute mechanical ventilation were recruited from the medical and surgical ICUs of an academic medical center in Northeast Ohio. Subject screening and recruitment were conducted from January 2011 to February 2012. Subjects were at least 50 years of age, required acute mechanical ventilation for less than 72 h, convalesced on an inpatient unit, made his/her own healthcare decisions, and did not have AD documentation in their medical records at ICU discharge. Hospitalized adults were excluded if they were unable to read and speak English, had severe vision or hearing impairment, or received inpatient palliative and/or hospice care.

Instrumentation

Acceptability

Acceptability of the educational interventions was measured by the single item question, "How helpful were the materials in assisting you to better understand advance directives"? This item was constructed as Likert scale from zero (not at all helpful) to 10 (extremely helpful). A mean score and standard deviation for the item was calculated to quantify acceptability.

Efficacy

Efficacy was assessed by a single item, "If given the chance today, would you complete a written advance directive"? with "do not intend to complete an AD" or "intend to complete an AD" as the responses. Immediately




Figure 1 Overview of subject flow through the study protocol.

upon enrollment, participants received the intervention. Thus, this question was administered posttest only to reduce a potential threat to internal validity, specifically a testing effect, from repeated exposures to this single item (Shadish, Cook, & Campbell, 2002).

Covariates

Based on prior research examining the effects of interventions to enhance AD completion rates (Alano et al., 2010; Salmond & Estrellam, 2005; Tamura et al., 2010), age, race, and level of preexisting knowledge of ADs have been shown to influence AD completion and were selected a priori as covariates. To capture age and race, each subject self-reported his or her age and racial identity. The Advance Directive Survey (ADS) was used to assess knowledge of ADs. For this survey, subjects endorse a response for each of the five items provided in a multiple-choice format. Correct answers are scored as 1, and incorrect answers are scored as 0. A total score, ranging from 0 to 5, was calculated by adding the number of correct re-

sponses. A higher ADS scores indicate that the subject has greater knowledge on ADs. This questionnaire has established construct validity in a sample of hospitalized male veterans (Johnson-Greene et al., 1996). In this sample of hospitalized adults, the Kuder Richardson (KR)-21 formula coefficient was 0.42, which indicates an adequate internal reliability and consistency for a dichotomously scored scale consisting of less than 10 items.

Procedures

Subject recruitment and screening

Institutional review board approval from the academic medical center was obtained prior to the conduct of this clinical research. Each weekday, the research assistant screened critically ill patients for study eligibility by review of the mechanical ventilation records and progress notes. All eligible critically ill patients transferred from the ICU to an inpatient unit were approached for study participation within 24–48 h after their admission to an inpatient unit.

Data collection

Data were collected by structured interviews at two time points, at baseline (24–48 h after transfer from an ICU) and 24 h following baseline (T2). Each interview was 30–40 min in duration. Clinical and demographic data were obtained from the patient's electronic and physical medical records and assessment of AD knowledge were collected at baseline. Data on intention to complete an AD and the acceptability of the intervention were collected at T2.

Data analysis

Data were analyzed using the Statistical Package for Social Sciences (SPSS, version 20; Somers, NY). Descriptive statistics, chi-square analyses, independent samples *t*-tests, and a logistic regression for multiple predictor variables were employed to evaluate acceptability, and efficacy of the MDS intervention compared to the AHA's brochure, *Putting It In Writing*. The criterion for statistical significance for all statistical tests included in this report was $p \leq .05$.

Acceptability

To determine the acceptability of the two educational interventions, descriptive statistics (means and standard deviations) of the single item measure, "How helpful were the materials in assisting you to better understand advance directives?" were generated, and an independent samples *t*-test was performed to determine if acceptability differed by group assignment.

Efficacy

A hierarchical logistic regression with multiple predictors was performed to assess the effects of group assignment on the outcome variable, intention to complete an AD, while adjusting for the potential influences of covariates (age, race, and AD knowledge). To determine the unique contribution of the covariates and the independent variable (group assignment), the covariates were simultaneously entered into model and the independent variable was subsequently entered.

Results

Sample characteristics

Table 1 shows the clinical and demographic characteristics of subjects by group assignment ($n = 49$). This convenience sample of hospitalized adults had a mean age of 63 ($SD = 9.5$) years. Most subjects were white

Table 1 Clinical and demographic characteristics of subjects by group assignment

Variable	Control ($n = 30$)		Experimental ($n = 19$)		<i>t</i>
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	
Age (years)	65.4	10.8	60.0	6.0	2.2*
ICU LOS (days)	2.7	2.4	4.0	5.9	-0.97
Hospital LOS (days)	10.2	6.9	13.3	9.3	-1.3
Charlson comorbidity index	2.7	6.8	3.1	4.2	-0.25
	<i>n</i>	%	<i>n</i>	%	χ^2
Gender: male	19	63	10	53	0.55
Race: white	18	60.0	13	68.4	0.35
Marital status: married	17	56.7	7	36.8	1.8
Annual income					
\$20,000 or less	7	25.9	4	21.1	0.18
\$20,001–49,999	10	37.0	7	36.8	
\$50,000 or more	10	37.0	8	42.1	

Note. ICU LOS, intensive care unit length of stay; hospital LOS, hospital length of stay.

* $p < .05$.

(63%), male (59%), transferred from the surgical ICU (84%), and experienced a mean ICU and hospital lengths of stay of 3.2 ($SD = 4.1$) days and 11.4 ($SD = 8.0$) days, respectively. There was a significant difference in age by group assignment. On average, the subjects exposed to the experimental condition were younger ($M = 60.0$, $SD = 6.0$) compared to subjects in the control group ($M = 65.4$, $SD = 10.8$, $t = 2.2$, $p < .05$). At baseline, AD knowledge scores did not differ by group assignment ($t = 0.85$, $p = .40$).

Acceptability

At T2, subjects ($n = 44$) completed a single item measure to assess the helpfulness of the educational resources delivered by each intervention. On average, subjects who were assigned to the MDS intervention ($M = 8.11$, $SD = 3.84$) found the educational content more helpful in improving their understanding of ADs compared to subjects exposed to the information-only control group ($M = 5.80$, $SD = 3.84$, $t = -2.67$, $p = .03$). The post hoc effect size was large ($d = 0.68$) and the observed power ($1 - \beta$) was 0.71 (one-tailed).

Efficacy

In order to examine the influence of MDS intervention on the outcome variable, intention to complete an AD, we conducted a hierarchical logistic regression analysis. In the initial step of the regression model, intention to complete an AD decision was regressed on covariates

Table 2 Hierarchical logistic regression of intention to complete an AD decision on age, race, AD knowledge, and group assignment ($n = 41$)

	Predictor	β	SE β	Exp β	p
Step 1	Age (0 = 50–64 years, 1 = ≥ 65 years)	–0.306	0.749	0.736	.736
	Race (0 = <i>white</i> , 1 = <i>non-white</i>)	1.091	0.755	2.98	.678
	AD knowledge	–0.658	0.315	0.518	.279
Step 2	Age	0.775	0.930	2.171	.405
	Race	1.984	1.006	7.275	.049*
	AD knowledge	–0.918	0.448	0.399	.040*
	Group assignment (0 = <i>control</i> , 1 = <i>experimental</i>)	3.206	1.608	24.690	.004**

Note. AD, advance directive; Dependent variable: intention to complete an AD.

Model statistics for step 1: $R^2 = 0.151$ (Cox & Snell), 0.203 (Nagelkerke). Model $\chi^2 = 6.697$, $df = 3$, $p = .082$. Hosmer & Lemeshow test ($\chi^2 = 12.408$, $df = 8$, $p = .137$).

Model statistics for step 2: $\Delta R^2 = 0.221$ (Cox & Snell), 0.298 (Nagelkerke). $R^2 = 0.372$ (Cox & Snell), 0.501 (Nagelkerke). Model $\chi^2 = 19.074$, $df = 4$, $p < .05$. Hosmer & Lemeshow test ($\chi^2 = 2.748$, $df = 7$, $p = .907$).

Overall correct classification was 79% for this model.

* $p < .05$, ** $p < .01$.

known to affect AD decision making (age, race, and AD knowledge), which did not achieve statistical significance. In the final step, the independent variable, group assignment, was added to evaluate its contribution to the model. As seen in Table 2, with study group assignment included in the model, the explained variance, Nagelkerke R^2 , was 0.501 and the overall correct classification of 79%. Based on this final model, subjects exposed to the MDS intervention were 24.7 times more likely to intend to complete an AD compared to those who were assigned to the control condition. In addition to exposure to the MDS intervention, having an effect on the intention to complete an AD, subjects who were white (OR = 7.275, $p = .049$) and having lower AD knowledge scores (OR = 0.399, $p = .040$) were found to be more likely to intend to complete an AD after controlling for age and group assignment.

Discussion

The results of this comparative quasi-experimental study establish the preliminary evidence for the acceptability and efficacy of the MDS intervention among hospitalized adults recovering from a critical illness. In the present study, our two educational interventions helped to increase subjects understanding of ADs. In particular, subjects assigned to the experimental condition found the interactive MDS intervention significantly more helpful compared to the educational brochure. Notably, this study confirms a statistically and clinically significant difference by group assignment in the proportion of subjects who intended to complete an AD. There was a significant difference in the proportion of subjects exposed to experimental condition who were more inclined to complete an AD compared to those who received the information only. Thus, these preliminary results are quite promis-

ing regarding the acceptability and efficacy of the MDS intervention. Our results using a single dose electronic intervention demonstrate similar efficacy as studies that have used combined approaches, such as counseling with education, to improve rates of AD completion (Heffner, Fahy, Hilling, & Barbieri, 1997; Payne, Prentice-Dunn, & Allen, 2009; Song et al., 2010). In addition to establishing the influence of the MDS intervention on intention to complete an AD decision, we found two significant predictors. Lower AD knowledge scores and individuals who identified their race as white were more likely to have the intention to complete an AD. Our findings regarding the influence of race and AD knowledge are consistent with the work of Sulmasy, Song, Marx, and Mitchell (1996) and Alano et al. (2010).

Limitations

There are several limitations that affect the internal and external validity of this research. First, the research design was quasi-experimental and subjects were not randomly assigned to the study conditions, but rather enrolled into the study sequentially. Second, the primary outcome, intention to complete an AD, was only captured after the subject was exposed to a study condition. This posttest-only design precluded statistically controlling for within-subject variation across time, but does reduce the threat of testing. Third, the research design did not account for a novelty effect of our MDS intervention, and future research is recommended to prospectively compare the efficacy of two electronic educational interventions. Lastly, the sample is small and consists of clinical and demographic characteristics that limit the external generalizability of the results to hospitalized adults who are white males with moderate levels of education and socioeconomic status.

Implications for practice

This pilot study established the acceptability and efficacy among hospitalized patients recovering from a critical illness. Educational interventions have been tested and efficacious, but have not considered the timing of intervention exposure which could impact receptiveness to establishing an AD and subsequently, the efficacy of intervention. In clinical practice, advance practice nurses seize naturally occurring opportunities, teachable moments, to provide education and foster health promotion behaviors. Yet, discussions about ADs are routinely omitted from the clinical management of acutely ill patients in hospital settings before discharge. This study provides evidence for a promising computer-mediated intervention to promote completion of an AD, while capitalizing on the use of technology to minimize the workload burden of registered nurses and advanced practice nurses. This technology allows patients to access educational materials and provides patients with a self-customized experience that promotes reflection, leveraging the teachable moment that may be embedded in the experience of critical illness. Future research should evaluate the essential components of the intervention, determine the mechanism by which the intervention works to increase intention to complete an AD, and examine the influence of the technology on the behavior of AD completion.

Acknowledgments

This publication was made possible by funding from Case Western Reserve University, Center on Aging and Health and the following grant awards: 2KLTR000440-06, L3ORR033212, and L30MH091738 from the National Center for Advancing Translation Science (NCATS), the former National Center for Research Resources (NCRR), and the National Institute of Mental Health (NIMH), components of the National Institutes of Health (NIH). Its contents are solely the responsibility of the authors and do not necessarily represent the official view of the NCATS, NCRR, NIMH, or the NIH.

References

- Alano, G., Pekmezaris, R., Tai, J., Hussain, M., Jeune, J., Louis, B., ... Wolf-Klein, G. (2010). Factors influencing older adults to complete advance directives. *Palliative and Supportive Care*, *8*(3), 267–275.
- Boudreaux, E., Bock, B., & O'Hea, E. (2012). When an event sparks behavior change: An introduction to the sentinel event method of dynamic model building and its application to emergency medicine. *Academic Emergency Medicine*, *19*, 329–335.
- Boudreaux, E., Moon, S., Baumann, B., Camargo, C., O'Hea, E., & Ziedonis, D. (2010). Intentions to quit smoking: Causal attribution, perceived illness severity, and event-related fear during an acute health event. *Annals of Behavioral Medicine*, *40*, 350–355.
- Cambhi, S., Mercado, A., Morrison, S., Du, Q., Platt, D., August, G., & Nelson, J. (2009). Deciding in the dark: Advance directives and continuation of treatment in chronic critical illness. *Critical Care Medicine*, *37*(3), 919–925.
- Clark, B., & Moss, M. (2011). Secondary prevention in the intensive care unit: Does ICU admission represent a "teachable moment"? *Critical Care Medicine*, *39*(6), 1500–1506.
- Cohen, D., Clark, E., Lawson, P., Casucci, B., & Flocke, S. (2011). Identifying teachable moments for health behavior counseling in primary care. *Patient Education and Counseling*, *85*(2), e8–e15.
- Cox, C., & Carson, S. (2007). An economic evaluation of prolonged mechanical ventilation. *Critical Care Medicine*, *35*(8), 1918–1927.
- Cox, C., Carson, S., & Lindquist, J. (2007). Differences in one year health outcomes and resource utilization by definition of prolonged mechanical ventilation: A prospective cohort study. *Critical Care Medicine*, *11*(1), 120–131.
- Cox, C., Martinu, T., Sathy, S., Clay, A., Chia, J., Gray, A., & Tulskey, J. (2009). Expectations and outcomes of prolonged mechanical ventilation. *Critical Care Medicine*, *37*(11), 2888–2894. doi: 10.1097/CCM.0b013e3181ab86ed
- Daly, B., & Douglas, S. (2005). Trial of a disease management program to reduce hospital readmissions of the chronically critically ill. *Chest*, *128*(2), 507–517.
- Douglas, S., & Daly, B. (2001). Hospital readmission among long-term ventilator patients. *Chest*, *120*(4), 1278–1286.
- Douglas, S., Daly, B., Gordon, N., & Brennan, P. (2002). Survival and quality of life: Short-term versus long-term ventilator patients. *Critical Care Medicine*, *30*(12), 2655–2662.
- Douglas, S., Daly, B., Kelley, C., O'Toole, E., & Montenegro, H. (2007). Chronically critically ill patients: Health-related quality of life and resource use after a disease management intervention. *Journal of Critical Care*, *16*(5), 447–457.
- Flocke, S., Antognoli, E., Step, M., Marsh, S., Parran, T., & Mason, J. (2012). A teachable moment communication process for smoking cessation talk: Description of a group randomized clinician-focused intervention. *BMC Health Services Research*, *12*(109), 1–13. doi: 10.1186/1472-6963-12-109
- Floyd, A., Steffens, R., Pavlik, E., & Andrykowski, M. (2011). Receipt of a false positive test result during routine screening for ovarian cancer: A teachable moment? *Journal of Clinical Psychology in Medical Settings*, *18*(1), 70–77.
- Heffner, J. E., Fahy, B., Hilling, L., & Barbieri, C. (1997). Outcomes of advance directive education of pulmonary rehabilitation patients. *American Journal of Respiratory and Critical Care Medicine*, *155*(3), 1055–1059.
- Johnson-Greene, D., Anderson, C., Adams, K., Buchtel, H., Miller, T., & Dehring, M. (1996). The psychologist's role in assessing and facilitating patients' knowledge of advance directives in medical settings: A preliminary investigation. *Journal of Clinical Psychology in Medical Settings*, *3*(2), 103–113.
- Kass-Bartelmes, B., Hughes, R., & Rutherford, M. (2003). Advance care planning: Preferences for care at end of life. *Research in Action* (12), 1–20. Retrieved from <http://www.scribd.com/doc/47966710/AHRQ-Research-In-Action-Advance-Care-Planning>
- Payne, K. L., Prentice-Dunn, S., & Allen, R. S. (2009). A comparison of two interventions to increase completion of advance directives. *Clinical Gerontologist*, *33*(1), 49–61.
- Salmond, S., & Estrellam, D. (2005). Attitudes toward advance directives and advance directive completion rates. *Orthopaedic Nursing*, *24*(2), 117–127.
- Shadish, W., Cook, T., & Campbell, D. (2002). *Experimental and quasi-experimental designs for generalized causal inference*. Boston, MA: Houghton Mifflin. p. 55.
- Sommers, M., Lyons, M., Bohn, M., Ribak, C., & Fargo, J. (2013). Health-compromising behaviours among young adults in the urban emergency department: Opportunity for a teachable moment. *Clinical*

- Nursing Research*. Epub Ahead of Print; 1–25 doi: 10.1177/1054773812471971
- Song, J., Ratner, E., Bartels, D., Ulvestad, N., Petroskas, D., West, M., . . . Gelberg, L. (2010). Effect of an end-of-life planning intervention on the completion of advance directives in homeless persons: A randomized trial. *Annals of Internal Medicine*, 153(2), 76–84.
- Sulmasy, D., Song, K., Marx, E., & Mitchell, J. (1996). Strategies to promote the use of advance directives in a residency outpatient practice. *Journal of General Internal Medicine*, 11(11), 657–663.
- Tamura, M., Goldstien, M., & Perez-Stable, E. (2010). Preferences for dialysis withdrawal and engagement in advance care planning within a diverse sample of dialysis patients. *Nephrology Dialysis Transplantation*, 25(1), 237–242.

Copyright of Journal of the American Association of Nurse Practitioners is the property of Wiley-Blackwell and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.